

ON THE CLASSES OF RAPIDLY VARYING FUNCTIONS

Slavko Simić

Communicated by Stevan Pilipović

ABSTRACT. The classes KR_∞ , MR_∞ , R_∞ of rapidly varying functions are natural extensions of Karamata's concept of regular variation. In [2] we introduced a new class K of perfect Karamata's kernels and its subclasses Θ and Σ . In this paper we study inclusion properties of these classes and, among other results, we prove $KR_\infty \subset MR_\infty \subset \Sigma \subset \Theta \subset K$.

Introduction

We begin with some definitions from Karamata's theory. A positive measurable function ℓ is *slowly varying in Karamata's sense* if $\ell(\lambda x) \sim \ell(x)$ ($x \rightarrow \infty$), for each $\lambda > 0$. Functions of the form $x^\rho \ell(x)$, $\rho \in \mathbb{R}$ are *regularly varying* with index ρ [1]. For a positive measurable function f , define \tilde{f} by $\tilde{f}(x) := \frac{f(x)}{\int_1^x f(t)/t dt}$. It is well known [1], that $\tilde{f}(x) \rightarrow \rho$, $0 < \rho < \infty$ ($x \rightarrow \infty$), if and only if f is regularly varying function in Karamata's sense with index ρ .

From there it follows an extension to the class Θ of rapidly varying functions. In [2] we gave the following definition.

DEFINITION 1. A positive measurable function p belongs to the class Θ if and only if $\tilde{p}(x) \rightarrow \infty$ ($x \rightarrow \infty$).

There is no representation form for the class Θ since its structure is ambiguous. For example, we showed in [2] that it is not closed under multiplication.

DEFINITION 2. Let Σ denote the maximal subclass of Θ which is closed under multiplication. Then Σ consists of all positive measurable functions s such that $s^2 \in \Theta$ [2, Theorem 1].

We also introduced the class K of *perfect Karamata's kernels*.

DEFINITION 3. A positive measurable kernel $C(\cdot)$ belongs to the class K if the asymptotic relation $\int_1^x f(t)C(t) dt \sim f(x) \int_1^x C(t) dt$ ($x \rightarrow \infty$), takes place for every regularly varying function $f(\cdot)$ of arbitrary index.

It is proved in [2] that a necessary and sufficient condition for $C \in K$ is

$$(1) \quad \int_1^x C(t) dt \in \Theta.$$

Strict inclusion [2],

$$(2) \quad \Sigma \subset \Theta \subset K,$$

takes place in the sense that Θ/Σ and K/Θ are not empty.

From the property of regularly varying function f with index ρ , $\forall \lambda > 0$, $f(\lambda x)/f(x) \rightarrow \lambda^\rho$ ($x \rightarrow \infty$), a natural extension to the class R_∞ arises.

DEFINITION 4. [1, p. 83] A positive measurable function f belongs to the class R_∞ if $f(\lambda x)/f(x) \rightarrow \infty$ ($x \rightarrow \infty$), for each $\lambda > 1$.

Subclasses of R_∞ are KR_∞ and MR_∞ .

DEFINITION 5. [1, p. 85] Let f be positive and measurable. Then

$$(i) \quad f \in KR_\infty \text{ if and only if } \liminf_{x \rightarrow \infty} \inf_{\lambda \geq 1} \frac{f(\lambda x)}{\lambda^c f(x)} = 1 \text{ for every } c \in R,$$

$$(ii) \quad f \in MR_\infty \text{ if and only if } \liminf_{x \rightarrow \infty} \inf_{\lambda \geq 1} \frac{f(\lambda x)}{\lambda^d f(x)} > 0 \text{ for every } d \in R,$$

There is strict inclusion [1, p. 83]

$$(3) \quad KR_\infty \subset MR_\infty \subset R_\infty.$$

We shall investigate intermediate inclusion properties of the classes KR_∞ , MR_∞ , R_∞ and Σ , Θ , K apart from (2) and (3).

Results

In all cases there is a strict inclusion property between the classes of rapidly varying functions mentioned above, except in the following one.

PROPOSITION 1. *The classes R_∞ and Θ are incomparable i.e., they have not an inclusion property.*

Because of the assertion above, there are two inclusion chains. The first one is

PROPOSITION 2. *An extension of (3) is the following*

$$KR_\infty \subset MR_\infty \subset R_\infty \subset K.$$

The second one is

PROPOSITION 3. *An extension of (2) is the following*

$$KR_\infty \subset MR_\infty \subset \Sigma \subset \Theta \subset K.$$

Therefore the class K includes all known classes of rapidly varying functions in Karamata's sense.

Proofs

PROOF OF PROPOSITION 1. In order to prove that the classes R_∞ and Θ are incomparable, we have to find some positive measurable functions f and g such that $f \in R_\infty$ but $f \notin \Theta$ and $g \in \Theta$ but $g \notin R_\infty$. \square

An example of f is the next one. Let $f(x) := xe^x$ except at the points $x = e^n$, $n \in N$, where we put $f(e^n) := e^{e^n - n}$. Now, using Definition 4, it is easy to verify that $f \in R_\infty$. But

$$\tilde{f}(e^n) = e^{e^n - n} / \int_1^{e^n} e^t dt \rightarrow 0 \quad (n \rightarrow \infty).$$

Hence $\liminf_{x \rightarrow \infty} \tilde{f}(x) = 0$, and $f \notin \Theta$.

An example of g is the following: denote by (p_n) , $n \in N$ the sequence of primes and let $g(x) := xe^x$ except at the points $x = p_n$ where $g(p_n) := p_n e^{2p_n}$. Since $g(x) \geq xe^x$ for $x \geq 1$, we get

$$\tilde{g}(x) \geq xe^x / \int_1^x e^t dt \rightarrow \infty \quad (x \rightarrow \infty);$$

hence $g \in \Theta$. But $\liminf_{x \rightarrow \infty} \frac{g(2x)}{g(x)} = 2$, i.e., $g \notin R_\infty$.

In order to prove Proposition 2, taking into account (3), we just have to prove that then $f \in K$ whenever $f \in R_\infty$. For this we need the following two lemmas.

LEMMA 1. *If $f \in R_\infty$, then $\int_1^x f(t) dt \in KR_\infty$.*

PROOF. Denote by $F(x) := \int_1^x f(t) dt$, and let $f \in R_\infty$. Since, for fixed $\lambda > 1$, $f(\lambda t)/f(t) \rightarrow \infty$ ($t \rightarrow \infty$) (Definition 4), for any $A > 0$ we can find t_0 such that $f(\lambda t) > Af(t)$ for $t > t_0 > 1$. Now, for sufficiently large x , we get

$$\frac{F(\lambda x)}{F(x)} = \frac{F(t_0) + \int_{t_0}^{\lambda x} f(t) dt}{F(t_0) + \int_{t_0}^x f(t) dt} > \frac{F(t_0) + \lambda \int_{t_0}^x f(\lambda t) dt}{F(t_0) + \int_{t_0}^x f(t) dt} > \frac{F(t_0) + \lambda A \int_{t_0}^x f(t) dt}{F(t_0) + \int_{t_0}^x f(t) dt} > A,$$

since $f(t) \rightarrow \infty$ ($t \rightarrow \infty$). Since A can be arbitrary large, we conclude that $F(x) \in R_\infty$. But $F(x)$ is also monotone increasing, hence [1, p. 85] $F \in KR_\infty$. \square

LEMMA 2. *If $g \in MR_\infty$ then $g \in \Theta$. Hence $MR_\infty \subset \Theta$.*

This lemma is proved in [1, p. 104].

PROOF OF PROPOSITION 2. Since $KR_\infty \subset MR_\infty(3)$, from the above lemmas we get $F(x) = \int_1^x f(t) dt \in \Theta$. Applying (1), we obtain $f \in K$. Hence $R_\infty \subseteq K$.

To prove strict inclusion we shall consider a function f_1 defined as: $f_1(x) := e^x$ except at the points $x = 2^n$, $n \in N$ where we put $f_1(2^n) := 2^n$. Then, clearly $\int_1^x f_1(t) dt \in \Theta$; hence by (1), $f_1 \in K$. Yet

$$\liminf_{x \rightarrow \infty} \frac{f_1(2x)}{f_1(x)} = 2,$$

hence $f_1 \notin R_\infty$. \square

PROOF OF PROPOSITION 3. From (2) and (3) follows that we have to prove that MR_∞ is a proper subclass of Σ . Applying Lemma 2 we obtain $KR_\infty \subset MR_\infty \subset \Theta$. But from Definition 5 evidently follows that if $f \in MR_\infty$ then also $f^2 \in MR_\infty \subset \Theta$. Hence, according to Definition 2, $MR_\infty \subseteq \Sigma$.

To prove that the class MR_∞ is a proper subclass of Σ , we shall consider the following example. Let $f(x) := \sqrt{\log x} \exp(\log^2 x)$, $x \geq 1$ except on intervals of the form $(\exp(n - 1/n), \exp n]$, $n \in \mathbb{N}$, where we put $f(x) := \sqrt{\log x} \exp(\log^2 x) / \sqrt[4]{n}$. We have to prove that $f \in \Sigma$, i.e., $f^2 \in \Theta$. In order to make calculations simpler, let us change the scale: $x \rightarrow \exp x$. In terms of $h(x) := f(e^x)$, we obtain

$$\tilde{f}^2(e^x) = \frac{f^2(e^x)}{\int_1^{e^x} f^2(t)/t dt} = \frac{h^2(x)}{\int_0^x h^2(t) dt}.$$

Then for $x > 0$,

$$\int_0^x h^2(t) dt < \int_0^x t e^{2t^2} dt < e^{2x^2}.$$

Hence for $x \notin \bigcup_{n=1}^{\infty} (n - 1/n, n]$,

$$\tilde{f}^2(e^x) = \frac{h^2(x)}{\int_0^x h^2(t) dt} > \frac{x e^{2x^2}}{e^{2x^2}} \rightarrow \infty \quad (x \rightarrow \infty).$$

If $x \in (n - 1/n, n]$ we obtain

$$\int_0^x h^2(t) dt = \int_0^{n-1/n} h^2(t) dt + \int_{n-1/n}^x h^2(t) dt < \exp(2(n - 1/n)^2) + \frac{e^{2x^2}}{\sqrt{n}}.$$

Hence

$$\begin{aligned} \tilde{f}^2(e^x) &> \frac{x e^{2x^2} / \sqrt{n}}{\exp(2(n - 1/n)^2) + e^{2x^2} / \sqrt{n}} = \frac{x}{1 + \sqrt{n} \exp(2(n - 1/n)^2 - 2x^2)} \\ &> \frac{n - 1/n}{\sqrt{n} + 1} \rightarrow \infty \quad (x \rightarrow \infty). \end{aligned}$$

Therefore we proved that $f^2 \in \Theta$. By Definition 2 this means that $f \in \Sigma$. Yet

$$\inf_{t \geq 0} \frac{h(n - 1/n + t)}{h(n - 1/n)} = \frac{1}{\sqrt[4]{n}}.$$

Hence

$$\liminf_{x \rightarrow \infty} \inf_{\lambda \geq 1} \frac{f(\lambda x)}{f(x)} = 0,$$

i.e., by Definition 5(i), $f \notin MR_\infty$. This yields the strict inclusion $MR_\infty \subset \Sigma$. Therefore Proposition 3 is proved. \square

REMARK 1. From Definition 3, it follows that if a function f is in the class K , it is still in K if changed in a denumerable number of points.

This remark is e.g., useful if one wants to verify that $\Theta \neq K$. Suppose $f_1 \in K$ is arbitrary. Define $f_0(n) = \int_1^n f_1(s)s^{-1}ds$ for $n = 1, 2, \dots$ and $f_0 = f_1$ elsewhere. Then $f_0 \in K$, $f_0 \notin \Theta$.

A similar remark applies to the proof of Proposition 2. The definition of $f_1 := e^x$ is irrelevant. Take $f_1 \in K$ arbitrary. Then define $f_0(2^n) = 2^n$ for $n \in \mathbb{N}$ and $f_0 = f_1$ elsewhere. Then $f_0 \notin R_\infty$ and $F_0 \in K$.

Since there is no representation (except for KR_∞) of rapidly varying functions, any information about it is welcomed. We can provide here such a one.

COROLLARY 1. *If $f \in R_\infty$, then*

$$\int_1^x f(t) dt = \exp\left(y(x) + z(x) + \int_1^x \frac{u(t)}{t} dt\right),$$

where $y(x)$ is non-decreasing and $z(x) \rightarrow 0$, $u(x) \rightarrow \infty$ ($x \rightarrow \infty$).

This result is a combination of Lemma 1 and well-known representation for the class KR_∞ [1, p. 86].

References

- [1] N. H. Bingham, C. M. Goldie, J. I. Teugels *Regular Variation*, Cambridge University Press, 1987.
- [2] S. Simić, *Integral kernels with regular variation property*, Publ. Inst. Math. Nouv. Sér. 72(86) (2002), 55–61.

Matematički institut SANU
Kneza Mihaila 35
11001 Beograd, p.p. 367
Serbia

ssimic@mi.sanu.ac.yu

(Received 06 12 2002)

(Revised 25 04 2003)