

**SOME CLASSES OF INTEGRAL GRAPHS
 WHICH BELONG TO THE CLASS $\overline{\alpha K_a \cup \beta K_{b,b}}$**

Mirko Lepović

Communicated by Slobodan Simić

ABSTRACT. Let G be a simple graph and let \overline{G} denote its complement. We say that G is integral if its spectrum consists of integral values. We have recently established a characterization of integral graphs which belong to the class $\overline{\alpha K_a \cup \beta K_{b,b}}$, where mG denotes the m -fold union of the graph G . In this work we investigate integral graphs from the class $\overline{\alpha K_a \cup \beta K_{b,b}}$ with $\overline{\lambda}_1 = a+b$, where $\overline{\lambda}_1$ is the largest eigenvalue of $\overline{\alpha K_a \cup \beta K_{b,b}}$.

In this work we consider only simple graphs. The spectrum of a simple graph G of order n contains the eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ of the ordinary adjacency matrix of G , and is denoted by $\sigma(G)$. A graph G is called integral if its spectrum $\sigma(G)$ consists only of integers [1].

An eigenvalue μ of G is main if and only if $\langle \mathbf{j}, \mathbf{Pj} \rangle = n \cos^2 \alpha > 0$, where \mathbf{j} is the main vector (with coordinates equal to 1) and \mathbf{P} is the orthogonal projection of the space \mathbb{R}^n onto the eigenspace $\mathcal{E}_A(\mu)$. The quantity $\beta = |\cos \alpha|$ is called the main angle of μ .

Let K_n and $K_{m,n}$ denote the complete graph and the complete bipartite graph, respectively. We have recently described all integral graphs which belong to the classes $\overline{\alpha K_a \cup \beta K_b}$, $\overline{\alpha K_{a,b}}$, $\overline{K_{a,a} \cup K_{b,b}}$ and $\overline{\alpha K_a \cup \beta K_{b,b}}$ (see [2], [3], [4] and [5], respectively), where \overline{G} and mG denote the complementary graph of G and the m -fold union of the graph G , respectively.

The characterization of integral graphs which is related to the class $\overline{\alpha K_a \cup \beta K_{b,b}}$ is reduced to the problem of finding the most general integral solution of the following Diophantine equation [5]

$$(1) \quad [(\alpha + 1)a + (2\beta - 1)b - 1]^2 - 4\alpha a(a - b - 1) = \delta^2.$$

In other words, $\overline{\alpha K_a \cup \beta K_{b,b}}$ is integral if and only if $(\alpha, \beta, a, b, \delta)$ represents a positive integral solution of the equation (1). We note that $\alpha K_a \cup \beta K_{b,b}$ is an

2000 *Mathematics Subject Classification:* Primary 05C50.

Key words and phrases: graph, eigenvalue, Diophantine equation, continued fractions.

integral graph with two main eigenvalues $\mu_a = a-1$ and $\mu_b = b$ for any $\alpha, \beta, a, b \in \mathbb{N}$ with $a \neq (b+1)$.

REMARK 1. We know that $\lambda_1 + \bar{\lambda}_1 \geq n-1$ for any graph G of order n with equality if and only if G is regular [1], where $\bar{\lambda}_1$ is the largest eigenvalue of \bar{G} . If $G = \alpha K_a \cup \beta K_{b,b}$ we obtain (i) $\bar{\lambda}_1 \geq 2b+1$ if $a > (b+1)$ and (ii) $\bar{\lambda}_1 \geq a+b$ if $a \leq b$.

In the sequel the symbol (m, n) denotes the greatest common divisor of integers m, n while $m \mid n$ means that m divides n .

THEOREM 1 (Lepović [5]). *If $\overline{\alpha K_a \cup \beta K_{b,b}}$ is integral then it belongs to one of the following classes of integral graphs:*

$$(2) \quad \left[\pm \frac{2kt}{\tau} x_0 + \frac{4mt}{\tau} z \right] K_a \cup \left[\pm \frac{2kt}{\tau} y_0 + \frac{a}{\tau} z \right] (2n-1) K_{b,b},$$

where (i) $a = \pm [2t + (2\ell - 1)(2n - 1)]k + (2\ell - 1)m + 1$ and $b = (2\ell - 1)m$;
(ii) $t, k, \ell, m, n \in \mathbb{N}$ such that $(m, 2n - 1) = 1$, $(2n - 1, 2t) = 1$ and $(2\ell - 1, 2t) = 1$;
(iii) $\tau = (a, 4mt)$ such that $\tau \mid 2kt$; (iv) x_0 and y_0 is a particular solution of the linear Diophantine equation $ax - (4mt)y = \tau$ and (v) $z \geq z_0$, where $z_0 = \min \mathbb{Z}$ such that $(\pm \frac{2kt}{\tau} x_0 + \frac{4mt}{\tau} z_0) \geq 1$ and $(\pm \frac{2kt}{\tau} y_0 + \frac{a}{\tau} z_0) \geq 1$;

$$(3) \quad \left[\pm \frac{(2t-1)k}{\tau} x_0 + \frac{2m(2t-1)}{\tau} z \right] K_a \cup \left[\pm \frac{(2t-1)k}{\tau} y_0 + \frac{a}{\tau} z \right] (2n-1) K_{b,b},$$

where (i) $a = \pm [(2t - 1) + (2\ell - 1)(2n - 1)]k + (2\ell - 1)m + 1$ and $b = (2\ell - 1)m$;
(ii) $t, k, \ell, m, n \in \mathbb{N}$ such that $(m, 2n - 1) = 1$, $(2n - 1, 2t - 1) = 1$ and $(2\ell - 1, 2t - 1) = 1$;
(iii) $\tau = (a, 2m(2t - 1))$ such that $\tau \mid (2t - 1)k$; (iv) x_0 and y_0 is a particular solution of the linear Diophantine equation $ax - 2m(2t - 1)y = \tau$ and (v) $z \geq z_0$ where $z_0 = \min \mathbb{Z}$ such that $(\pm \frac{(2t-1)k}{\tau} x_0 + \frac{2m(2t-1)}{\tau} z_0) \geq 1$ and $(\pm \frac{(2t-1)k}{\tau} y_0 + \frac{a}{\tau} z_0) \geq 1$;

$$(4) \quad \left[\pm \frac{(2t-1)k}{\tau} x_0 + \frac{(2t-1)m}{\tau} z \right] K_a \cup \left[\pm \frac{(2t-1)k}{\tau} y_0 + \frac{a}{\tau} z \right] n K_{b,b},$$

where (i) $a = \pm [(2t - 1) + 2\ell n]k + \ell m + 1$ and $b = \ell m$; (ii) $t, k, \ell, m, n \in \mathbb{N}$ such that $(m, n) = 1$, $(n, 2t - 1) = 1$ and $(\ell, 2t - 1) = 1$; (iii) $\tau = (a, (2t - 1)m)$ such that $\tau \mid (2t - 1)k$; (iv) x_0 and y_0 is a particular solution of the linear Diophantine equation $ax - (2t - 1)my = \tau$ and (v) $z \geq z_0$ where $z_0 = \min \mathbb{Z}$ with $(\pm \frac{(2t-1)k}{\tau} x_0 + \frac{(2t-1)m}{\tau} z_0) \geq 1$ and $(\pm \frac{(2t-1)k}{\tau} y_0 + \frac{a}{\tau} z_0) \geq 1$. In these classes the symbol ‘ \pm ’ is related to ‘+’ if $a > (b+1)$; and ‘ \pm ’ is related to ‘-’ if $a \leq b$.

If $\overline{\alpha K_a \cup \beta K_{b,b}}$ is an integral graph then it uniquely determines the parameters t, τ, k, ℓ, m, n . However, if x_0 and y_0 is obtained by using the EUCLID algorithm then a fixed integral graph $\overline{\alpha K_a \cup \beta K_{b,b}}$ also uniquely determines the parameters x_0, y_0, z_0, z (see [5]).

Using Theorem 1 we proved in [5] the following results: (i) if $\overline{\alpha K_a \cup \beta K_{b,b}}$ is integral with $\bar{\lambda}_1 = 2b+1$ and $a > (b+1)$ then it is $\overline{K_5 \cup K_{2,2}}$; (ii) if $\overline{\alpha K_a \cup \beta K_{b,b}}$

is integral with $\bar{\lambda}_1 = 2b + 1$ and $a \leq b$ then it belongs to the class of integral graphs $\overline{3K_t \cup K_{4t-2, 4t-2}}$, where $t \in \mathbb{N}$ and (iii) if $\overline{\alpha K_a \cup \beta K_{b,b}}$ is integral with $\bar{\lambda}_1 = a + b$ and $a \leq b$ then it is one of the following two integral graphs $\overline{K_2 \cup K_{6,6}}$ or $\overline{K_3 \cup K_{6,6}}$.

The characterization of integral graphs with $\bar{\lambda}_1 = a + b$ and $\lambda_1 = a - 1$ is reduced to the problem of finding the most general positive solution of the equation $x^2 - dy^2 = c$, where d is not a perfect square. It is based on the concept of continued fractions and some basic results which are related to $x^2 - dy^2 = c$ (see [6]).

Let a_0, a_1, \dots, a_n be a sequence of integers with $a_i > 0$ for $i \geq 1$. Then the term $[a_0; a_1, \dots, a_n] = [a_0; a_1, \dots, a_{n-1} + \frac{1}{a_n}]$ is called the simple continued fraction, where $[a_0; a_1] = a_0 + \frac{1}{a_1}$. If a_0, a_1, a_2, \dots is an infinite sequence of integers with $a_i > 0$ for $i \geq 1$, the expression $[a_0; a_1, a_2, \dots] = \lim_{n \rightarrow +\infty} [a_0; a_1, \dots, a_n]$ is called the infinite simple continued fraction. We say that $[a_0; a_1, \dots, a_{m-1}, \overline{a_m, \dots, a_{m+r-1}}]$ is an infinite simple continued fraction of periodic r if r is the least positive integer such that $a_{r+n} = a_n$ for any $n \geq m$.

Let a_0, a_1, \dots be a sequence of integers with $a_i > 0$ for $i \geq 1$. We then define two associated sequences $\{p_n\}$ and $\{q_n\}$ by $p_i = a_i p_{i-1} + p_{i-2}$ and $q_i = a_i q_{i-1} + q_{i-2}$ for $i \geq 0$, where $p_{-2} = 0$, $p_{-1} = 1$ and $q_{-2} = 1$, $q_{-1} = 0$. The rational number $\frac{p_n}{q_n} = [a_0; a_1, \dots, a_n]$ is called the n -th convergent to the infinite simple continued fraction.

Next, the general solution of the Pell equation $x^2 - dy^2 = 1$ is given in the form $x_i + y_i \sqrt{d} = (x_1 + y_1 \sqrt{d})^i$, where $x_1 + y_1 \sqrt{d}$ is its fundamental solution. We know that $x_1 + y_1 \sqrt{d} = p_{r-1} + q_{r-1} \sqrt{d}$ if r is even, and $x_1 + y_1 \sqrt{d} = p_{2r-1} + q_{2r-1} \sqrt{d}$ if r is odd, where r is the period length of \sqrt{d} . If $\rho_0 + \varphi_0 \sqrt{d}$ is a fundamental solution of the equation $x^2 - dy^2 = c$, then

$$\rho_i + \varphi_i \sqrt{d} = (\rho_0 + \varphi_0 \sqrt{d})(x_1 + y_1 \sqrt{d})^i$$

represents a class of solutions of $x^2 - dy^2 = c$. Using the last relation we easily find that $\rho_i = \rho_0 x_i + d \varphi_0 y_i$ and $\varphi_i = \varphi_0 x_i + \rho_0 y_i$ for any $i \geq 0$, understanding that $x_0 = 1$ and $y_0 = 0$. Besides, we have

$$(5) \quad \rho_i = \frac{\rho_0 + \varphi_0 \sqrt{d}}{2} (x_1 + y_1 \sqrt{d})^i + \frac{\rho_0 - \varphi_0 \sqrt{d}}{2} (x_1 - y_1 \sqrt{d})^i;$$

$$(6) \quad \varphi_i = \frac{\rho_0 + \varphi_0 \sqrt{d}}{2\sqrt{d}} (x_1 + y_1 \sqrt{d})^i - \frac{\rho_0 - \varphi_0 \sqrt{d}}{2\sqrt{d}} (x_1 - y_1 \sqrt{d})^i.$$

Finally, for any fundamental solution $\rho_0 + \varphi_0 \sqrt{d}$ of the equation $x^2 - dy^2 = c$, the following two relations are satisfied [6]

$$(7) \quad 0 \leq |\rho_0| \leq \sqrt{\frac{c(x_1 + 1)}{2}} \quad \text{and} \quad 0 \leq \varphi_0 \leq y_1 \sqrt{\frac{c}{2(x_1 + 1)}}.$$

Using the concept of continued fractions we proved in [5] that there is no integral graph from the class $\overline{\alpha K_a \cup (3\beta + 2)K_{b,b}}$ with $\bar{\lambda}_1 = a + b$ and $a > (b + 1)$ for any $\beta \in \mathbb{N}$. It is also observed that there is no integral graph from the class $\overline{\alpha K_a \cup \beta K_{b,b}}$ with $\bar{\lambda}_1 = a + b$ and $a > (b + 1)$ for $\beta = 1$.

The characterization of integral graphs with $\bar{\lambda}_1 = a + b$ and $\lambda_1 = a - 1$ is reduced to the problem of finding the most general positive integral solution of the following two Diophantine equations:

$$(8) \quad [8\eta(\eta\dot{n} - 1)m - k]^2 - [16\eta\dot{n}(\eta\dot{n} - 1) + 1]k^2 = 16\eta(\eta\dot{n} - 1),$$

where $\dot{n} = 2n - 1$ and $\beta = \eta\dot{n}$; (1.1) $a = (2\ell - 1)(2t - 1)$; (1.2) $b = (2\ell - 1)m$; (1.3) $(2\ell - 1) = 2\eta m + k$ and (1.4) $(2t - 1) = (2\ell - 1)\dot{n} - m$; and

$$(9) \quad [4\eta(\eta n - 1)m - k]^2 - [16\eta n(\eta n - 1) + 1]k^2 = 8\eta(\eta n - 1),$$

where $\beta = \eta n$ and (2.1) $a = (2t - 1)\ell$; (2.2) $b = \ell m$; (2.3) $\ell = \eta m + k$ and (2.4) $(2t - 1) = 2\ell n - m$ (see [5]).

Further, let $x = 8\eta(\eta\dot{n} - 1)m - k$ and let $y = k$. Let $d = 16\eta\dot{n}(\eta\dot{n} - 1) + 1$ and let $\rho_0 + \varphi_0\sqrt{d}$ be a fundamental solution of $x^2 - dy^2 = 16\eta(\eta\dot{n} - 1)$. Then $k = \varphi_i$ and $m = \frac{\rho_i + \varphi_i}{8\eta(\eta\dot{n} - 1)}$, understanding that $\rho_i + \varphi_i\sqrt{d}$ is the i -th solution which belongs to the class with respect to $\rho_0 + \varphi_0\sqrt{d}$. It was proved in [5] that $8\eta(\eta\dot{n} - 1) \mid (\rho_i + \varphi_i)$ if and only if $8\eta(\eta\dot{n} - 1) \mid (\rho_0 + \varphi_0)$. Consequently, the most general integral solution of (8) is reduced to the positive fundamental solutions $\rho_0 + \varphi_0\sqrt{d}$ for which $8\eta(\eta\dot{n} - 1) \mid (\rho_0 + \varphi_0)$. Similarly, the most general integral solution of (9) is reduced to the positive fundamental solutions $\rho_0 + \varphi_0\sqrt{d}$ for which $4\eta(\eta n - 1) \mid (\rho_0 + \varphi_0)$.

We now proceed to establish a characterization of integral graphs $\overline{\alpha K_a \cup \beta K_{b,b}}$ with $\bar{\lambda}_1 = a + b$ and $a > (b + 1)$ for $\beta = 2, 3, 4$. We note first if $\overline{\alpha K_a \cup \beta K_{b,b}}$ is an integral graph with $\bar{\lambda}_1 = a + b$ and $\lambda_1 = a - 1$ then $(a + b) + (a - 1) \geq \alpha a + 2\beta b$ (see Remark1), which implies that $\alpha = 1$.

PROPOSITION 1. *If $\overline{\alpha K_a \cup 2K_{b,b}}$ is integral with $\bar{\lambda}_1 = a + b$ and $a > (b + 1)$ then it belongs to the following class of integral graphs*

$$\overline{K_{a_+ z_+^{2i} + a_- z_-^{2i} + \frac{1}{33}} \cup 2K_{b_+ z_+^{2i} + b_- z_-^{2i} + \frac{7}{33}, b_+ z_+^{2i} + b_- z_-^{2i} + \frac{7}{33}}}$$

where $z_{\pm} = 23 \pm 4\sqrt{33}$ and $i \geq 0$, $a_{\pm} = \frac{247 \pm 43\sqrt{33}}{33}$ and $b_{\pm} = \frac{46 \pm 8\sqrt{33}}{33}$.

PROOF. We shall first consider the general positive integral solution of the equation (8) for $\eta\dot{n} = 2$. Clearly, $\dot{n} = 1$ and $\eta = 2$. Then relation (8) is reduced to $x^2 - 33y^2 = 32$. Using a computer program¹ we obtain that $\sqrt{33} = [5; \overline{1, 2, 1, 10}]$ and $23 + 4\sqrt{33}$ is the fundamental solution of the equation $x^2 - 33y^2 = 1$. Since $\rho_0 \leq 19$ and $\varphi_0 \leq 3$ (see (7)), it is easy to verify that there is no fundamental solution of $x^2 - 33y^2 = 32$, which means that (8) does not generate any integral graph with $\beta = 2$.

Consider the general positive integral solution of the equation (9) for $\eta n = 2$. We shall distinguish the following two cases:

Case 1. ($n = 1$ and $\eta = 2$). Then (9) is reduced to (i) $x^2 - 33y^2 = 16$. We now find that $\rho_0 \leq 13$ and $\varphi_0 \leq 2$, and $4 + 0\sqrt{33}$ and $7 + \sqrt{33}$ are the fundamental

¹All the results given in Propositions1,2 and3 are obtained by using the program called DIOPHANTUS, written by the author in the programming language C.

solutions of (i). Since $8 \nmid (4 + 0)$ it follows that the class of solutions of (i) which corresponds to $4 + 0\sqrt{33}$ does not generate any integral graph with $\beta = 2$. Since $m = \frac{\rho_i + \varphi_i}{8}$ and $k = \varphi_i$, for the fundamental solution $7 + \sqrt{33}$, we obtain from (5) and (6) that

$$m = \frac{\sqrt{33} + 5}{2\sqrt{33}} \left(23 + 4\sqrt{33}\right)^i + \frac{\sqrt{33} - 5}{2\sqrt{33}} \left(23 - 4\sqrt{33}\right)^i;$$

$$k = \frac{7 + \sqrt{33}}{2\sqrt{33}} \left(23 + 4\sqrt{33}\right)^i - \frac{7 - \sqrt{33}}{2\sqrt{33}} \left(23 - 4\sqrt{33}\right)^i.$$

Further, making use of (2.1), (2.2), (2.3) and (2.4), from the previous relations we easily get

$$\ell = \frac{3\sqrt{33} + 17}{2\sqrt{33}} \left(23 + 4\sqrt{33}\right)^i + \frac{3\sqrt{33} - 17}{2\sqrt{33}} \left(23 - 4\sqrt{33}\right)^i;$$

$$t = \frac{5\sqrt{33} + 29}{2\sqrt{33}} \left(23 + 4\sqrt{33}\right)^i + \frac{5\sqrt{33} - 29}{2\sqrt{33}} \left(23 - 4\sqrt{33}\right)^i,$$

which provides the class of integral graphs represented in Proposition 1, understanding that $\dot{p} = 2p - 1$.

Case 2. ($n = 2$ and $\eta = 1$). Then (9) is reduced to (ii) $x^2 - 33y^2 = 8$. We now find that (iii) $\rho_0 \leq 9$ and $\varphi_0 \leq 1$. Using (iii) it is not difficult to show that there exists no fundamental solution of (ii), which completes the proof. \square

PROPOSITION 2. *If $\overline{\alpha K_a \cup 3K_{b,b}}$ is integral with $\bar{\lambda}_1 = a + b$ and $a > (b + 1)$ then it belongs to one of the following three classes of integral graphs:*

$$\overline{K_{a_+ z_{\pm}^{2i} + a_- z_{\pm}^{2i} + \frac{1}{97}} \cup 3K_{b_+ z_{\pm}^{2i} + b_- z_{\pm}^{2i} + \frac{11}{97}, b_+ z_{\pm}^{2i} + b_- z_{\pm}^{2i} + \frac{11}{97}}}$$

where $z_{\pm} = 62809633 \pm 6377352\sqrt{97}$ and $i \geq 0$; and

$$(1^0) \ a_{\pm} = \frac{6170687737 \pm 626538413\sqrt{97}}{97} \text{ and } b_{\pm} = \frac{1309509107 \pm 132960505\sqrt{97}}{194};$$

$$(2^0) \ a_{\pm} = \frac{5188723 \pm 526835\sqrt{97}}{194} \text{ and } b_{\pm} = \frac{550561 \pm 55901\sqrt{97}}{194} \text{ and}$$

$$(3^0) \ a_{\pm} = \frac{681412777 \pm 69186985\sqrt{97}}{194} \text{ and } b_{\pm} = \frac{72302819 \pm 7341239\sqrt{97}}{194}.$$

PROOF. We shall first consider the general positive integral solution of the equation (8) for $\eta\dot{n} = 3$.

Case 1.1 ($\dot{n} = 1$ and $\eta = 3$). Then (8) is reduced to (i) $x^2 - 97y^2 = 96$. We now have (ii) $\sqrt{97} = [9; 1, 5, 1, 1, 1, 1, 1, 5, 1, 18]$; (iii) $62809633 + 6377352\sqrt{97}$ is the fundamental solution of the equation $x^2 - 97y^2 = 1$ and (iv) $\rho_0 \leq 54907$ and $\varphi_0 \leq 5575$. According to (iv) we find that $22 + 2\sqrt{97}$; $463 + 47\sqrt{97}$; $2738 + 278\sqrt{97}$ and $49589 + 5035\sqrt{97}$ are the fundamental solutions of (i). Since $48 \nmid (22 + 2)$; $48 \nmid (463 + 47)$ and $48 \nmid (2738 + 278)$, these solutions do not generate any integral graph with $\beta = 3$.

Consequently, the general solution of (i) is reduced to the class which corresponds to the fundamental solution $49589 + 5035\sqrt{97}$. Since $m = \frac{\rho_i + \varphi_i}{48}$ and $k = \varphi_i$, using (iii) and (5), (6), we obtain

$$m = \left(\frac{1138\sqrt{97} + 11208}{2\sqrt{97}} \right) z_+^i + \left(\frac{1138\sqrt{97} - 11208}{2\sqrt{97}} \right) z_-^i;$$

$$k = \left(\frac{49589 + 5035\sqrt{97}}{2\sqrt{97}} \right) z_+^i - \left(\frac{49589 - 5035\sqrt{97}}{2\sqrt{97}} \right) z_-^i.$$

Next, making use of (1.1), (1.2), (1.3) and (1.4), by a straight-forward calculation, we get from the last relation that

$$\dot{\ell} = \left(\frac{11863\sqrt{97} + 116837}{2\sqrt{97}} \right) z_+^i + \left(\frac{11863\sqrt{97} - 116837}{2\sqrt{97}} \right) z_-^i;$$

$$\dot{i} = \left(\frac{10725\sqrt{97} + 105629}{2\sqrt{97}} \right) z_+^i + \left(\frac{10725\sqrt{97} - 105629}{2\sqrt{97}} \right) z_-^i,$$

which provides the class of integral graphs represented in Proposition 2 (1⁰).

Case 1.2 ($\dot{n} = 3$ and $\eta = 1$). Then (8) is reduced to (v) $x^2 - 97y^2 = 32$. According to (iii) and (7) we find that (vi) $\rho_0 \leq 31701$ and $\varphi_0 \leq 3218$. Using (vi) we get $138 + 14\sqrt{97}$ and $3063 + 311\sqrt{97}$ are the fundamental solutions of (v). Since $16 \nmid (138 + 14)$ and $16 \nmid (3063 + 311)$ it follows that (v) generates no integral graph with $\beta = 3$.

Consider the general positive integral solution of the equation (9) for $\eta n = 3$. We shall also distinguish the following two cases:

Case 2.1 ($n = 1$ and $\eta = 3$). Then (9) is reduced to (vii) $x^2 - 97y^2 = 48$. We now find that $\rho_0 \leq 38825$ and $\varphi_0 \leq 3942$; $40 + 4\sqrt{97}$, $719 + 73\sqrt{97}$ and $15965 + 1621\sqrt{97}$ are the fundamental solutions of (vii). Since $24 \nmid (40 + 4)$ and $24 \nmid (15965 + 1621)$ it remains to consider the fundamental solution $719 + 73\sqrt{97}$. Therefore, by an easy calculation we get $m = \left(\frac{33\sqrt{97} + 325}{2\sqrt{97}} \right) z_+^i + \left(\frac{33\sqrt{97} - 325}{2\sqrt{97}} \right) z_-^i$ and $k = \left(\frac{719 + 73\sqrt{97}}{2\sqrt{97}} \right) z_+^i - \left(\frac{719 - 73\sqrt{97}}{2\sqrt{97}} \right) z_-^i$, which yields $\ell = \left(\frac{86\sqrt{97} + 847}{\sqrt{97}} \right) z_+^i + \left(\frac{86\sqrt{97} - 847}{\sqrt{97}} \right) z_-^i$ and $\dot{i} = \left(\frac{311\sqrt{97} + 3063}{2\sqrt{97}} \right) z_+^i + \left(\frac{311\sqrt{97} - 3063}{2\sqrt{97}} \right) z_-^i$. So we get the class of integral graphs represented in Proposition 2 (2⁰).

Case 2.2 ($n = 3$ and $\eta = 1$). Then (9) is reduced to (viii) $x^2 - 97y^2 = 16$. We now find that $\rho_0 \leq 22416$ and $\varphi_0 \leq 2275$; $4 + 0\sqrt{97}$ and $4757 + 483\sqrt{97}$ are the fundamental solutions of (viii). Consequently, since $8 \nmid (4 + 0)$ and $8 \mid (4757 + 483)$ we obtain that $m = \left(\frac{655\sqrt{97} + 6451}{2\sqrt{97}} \right) z_+^i + \left(\frac{655\sqrt{97} - 6451}{2\sqrt{97}} \right) z_-^i$; $k = \left(\frac{4757 + 483\sqrt{97}}{2\sqrt{97}} \right) z_+^i - \left(\frac{4757 - 483\sqrt{97}}{2\sqrt{97}} \right) z_-^i$; $\ell = \left(\frac{569\sqrt{97} + 5604}{\sqrt{97}} \right) z_+^i + \left(\frac{569\sqrt{97} - 5604}{\sqrt{97}} \right) z_-^i$; $\dot{i} = \left(\frac{6173\sqrt{97} + 60797}{2\sqrt{97}} \right) z_+^i + \left(\frac{6173\sqrt{97} - 60797}{2\sqrt{97}} \right) z_-^i$, which provides the class represented in Proposition 2 (3⁰). \square

PROPOSITION 3. If $\overline{\alpha K_a \cup 4K_{b,b}}$ is integral with $\bar{\lambda}_1 = a + b$ and $a > (b + 1)$ then it belongs to one of the following three classes of integral graphs:

$$\overline{K_{a_+ z_+^{2i} + a_- z_-^{2i} + \frac{1}{193}} \cup 4K_{b_+ z_+^{2i} + b_- z_-^{2i} + \frac{15}{193}, b_+ z_+^{2i} + b_- z_-^{2i} + \frac{15}{193}}}$$

where $z_{\pm} = 6224323426849 \pm 448036604040\sqrt{193}$ and $i \geq 0$; and

$$(1^0) \ a_{\pm} = \frac{1209056824462393 \pm 87029814579823\sqrt{193}}{193} \text{ and}$$

$$b_{\pm} = \frac{179835915982455 \pm 12944872487449\sqrt{193}}{386};$$

$$(2^0) \ a_{\pm} = \frac{758972 \pm 54632\sqrt{193}}{193} \text{ and } b_{\pm} = \frac{56445 \pm 4063\sqrt{193}}{193};$$

$$(3^0) \ a_{\pm} = \frac{92695388006569 \pm 6672360030889\sqrt{193}}{386} \text{ and } b_{\pm} = \frac{6893786823015 \pm 496225633751\sqrt{193}}{386}.$$

PROOF. We shall first consider the general positive integral solution of the equation (8) for $\eta\dot{n} = 4$. Clearly, $\dot{n} = 1$ and $\eta = 4$. In this case (8) is reduced to (i) $x^2 - 193y^2 = 192$. We now have (ii) $\sqrt{193} = [13; \overline{1, 8, 3, 2, 1, 3, 3, 1, 2, 3, 8, 1, 26}]$; (iii) $6224323426849 + 448036604040\sqrt{193}$ is the fundamental solution of the Pell equation $x^2 - 193y^2 = 1$ and (iv) $\rho_0 \leq 24444530$ and $\varphi_0 \leq 1759555$. Using (iv) we find that $112 + 8\sqrt{193}$; $3362 + 242\sqrt{193}$; $87703 + 6313\sqrt{193}$; $871862 + 62758\sqrt{193}$ and $22743973 + 1637147\sqrt{193}$ are the fundamental solutions of (i). Since $96 \nmid (112 + 8)$; $96 \nmid (3362 + 242)$; $96 \nmid (87703 + 6313)$ and $96 \nmid (871862 + 62758)$, these solutions do not generate any integral graph with $\beta = 4$.

Thus, the general solution of (i) is reduced to the class which corresponds to the fundamental solution $22743973 + 1637147\sqrt{193}$. Making use of (iii) and (5), (6), we get implicitly that $m = (\frac{126985\sqrt{193} + 1764132}{\sqrt{193}})z_+^i + (\frac{126985\sqrt{193} - 1764132}{\sqrt{193}})z_-^i$ and $k = (\frac{22743973 + 1637147\sqrt{193}}{2\sqrt{193}})z_+^i - (\frac{22743973 - 1637147\sqrt{193}}{2\sqrt{193}})z_-^i$, which provides that $\ell = (\frac{3668907\sqrt{193} + 50970085}{2\sqrt{193}})z_+^i + (\frac{3668907\sqrt{193} - 50970085}{2\sqrt{193}})z_-^i$; $t = (\frac{3414937\sqrt{193} + 47441821}{2\sqrt{193}})z_+^i + (\frac{3414937\sqrt{193} - 47441821}{2\sqrt{193}})z_-^i$. So we arrive at the class of integral graphs represented in Proposition 3 (1^0).

Consider the general positive integral solution of the equation (9) for $\eta n = 4$. We shall distinguish the following three cases:

Case 1. ($n = 1$ and $\eta = 4$). Then (9) is reduced to (v) $x^2 - 193y^2 = 96$. We now find that $\rho_0 \leq 17284892$ and $\varphi_0 \leq 1244193$; $17 + \sqrt{193}$, $403 + 29\sqrt{193}$, $12142 + 874\sqrt{193}$ and $3148778 + 226654\sqrt{193}$ are the fundamental solutions of (v). Of course, since $48 \nmid (17 + 1)$; $48 \nmid (12142 + 874)$ and $48 \nmid (3148778 + 226654)$, these solutions generate no integral graph with $\beta = 4$. For $403 + 29\sqrt{193}$ we have $m = (\frac{9\sqrt{193} + 125}{2\sqrt{193}})z_+^i + (\frac{9\sqrt{193} - 125}{2\sqrt{193}})z_-^i$; $k = (\frac{403 + 29\sqrt{193}}{2\sqrt{193}})z_+^i - (\frac{403 - 29\sqrt{193}}{2\sqrt{193}})z_-^i$; $\ell = (\frac{65\sqrt{193} + 903}{2\sqrt{193}})z_+^i + (\frac{65\sqrt{193} - 903}{2\sqrt{193}})z_-^i$ and $t = (\frac{121\sqrt{193} + 1681}{2\sqrt{193}})z_+^i + (\frac{121\sqrt{193} - 1681}{2\sqrt{193}})z_-^i$, which provides the class of integral graphs represented in Proposition 3 (2^0).

Case 2. ($n = 2$ and $\eta = 2$). Then (9) is reduced to (vi) $x^2 - 193y^2 = 48$. We now find that $\rho_0 \leq 12222265$ and $\varphi_0 \leq 879777$; $56 + 4\sqrt{193}$, $1681 + 121\sqrt{193}$

and $435931 + 31379\sqrt{193}$ are the fundamental solutions of (vi). Consequently, since $24 \nmid (56 + 4)$, $24 \nmid (1681 + 121)$ and $24 \nmid (435931 + 31379)$, the equation (vi) does not generate any integral graph with $\beta = 4$.

Case 3. ($n = 4$ and $\eta = 1$). Then (9) is reduced to (vii) $x^2 - 193y^2 = 24$. We now find that $\rho_0 \leq 8642446$ and $\varphi_0 \leq 622096$; $6071 + 437\sqrt{193}$ and $1574389 + 113327\sqrt{193}$ are the fundamental solutions of (vii). Since $12 \nmid (6071 + 437)$ and $12 \mid (1574389 + 113327)$, we obtain for $1574389 + 113327\sqrt{193}$ that $m = \left(\frac{140643\sqrt{193} + 1953875}{2\sqrt{193}}\right)z_+^i + \left(\frac{140643\sqrt{193} - 1953875}{2\sqrt{193}}\right)z_-^i$ and $k = \left(\frac{1574389 + 113327\sqrt{193}}{2\sqrt{193}}\right)z_+^i - \left(\frac{1574389 - 113327\sqrt{193}}{2\sqrt{193}}\right)z_-^i$. In this way we obtain that $\ell = \left(\frac{126985\sqrt{193} + 1764132}{\sqrt{193}}\right)z_+^i + \left(\frac{126985\sqrt{193} - 1764132}{\sqrt{193}}\right)z_-^i$ and $t = \left(\frac{1891117\sqrt{193} + 26272237}{2\sqrt{193}}\right)z_+^i + \left(\frac{1891117\sqrt{193} - 26272237}{2\sqrt{193}}\right)z_-^i$. Using these relations we obtain Proposition 3 (3^0). \square

Table 1 contains the set of all integral graphs² from the class $\overline{\alpha K_a \cup \beta K_{b,b}}$, whose order 'o' does not exceed 30. In this table an integral graph is described by the parameters α, β, a, b and ones presented in the class of integral graphs in Theorem 1. The symbol 'i' denotes the identification number of the corresponding integral graph. In Table 1 (i) graphs with identification numbers 1, 2, ..., 18 belong to the classes represented by (2); (ii) graphs with identification numbers 19, 20, ..., 47 belong to the classes represented by (3); and (iii) graphs with $i = 48, 49, \dots, 70$ belong to the classes represented by (4). We note that there exist exactly 18, 29 and 23 non-isomorphic integral graphs from the classes described by (2), (3) and (4), respectively. In this table³ identification number 20 is related to the integral graph with the largest eigenvalue $\bar{\mu}_1 = 2b + 1$ and $a > (b + 1)$, while identification numbers 4, 19 and 44 are related to the integral graphs with $\bar{\mu}_1 = 2b + 1$ and $a \leq b$. In Table 1 there exists just one integral graph⁴ with $\bar{\mu}_1 = (a + b)$ and $a > (b + 1)$ and its identification number is 64 – the first next one has 12545 vertices. Identification numbers 24 and 50 are related to the integral graphs with $\bar{\mu}_1 = (a + b)$ and $a \leq b$.

There exist exactly 7556 non-isomorphic integral graphs which belong to the class $\overline{\alpha K_a \cup \beta K_{b,b}}$, whose order does not exceed 300. In particular, the total number of such integral graphs (obtained by using (2), (3) and (4)) is $(1433 + 888)$, $(1265 + 948)$ and $(1736 + 1286)$, respectively, where m and n in the expression $(m + n)$ are the numbers of integral graphs with $a > (b + 1)$ and $a \leq b$, respectively. Table 2 contains a distribution of those graphs with respect to their orders. In Table 2 the number n in the symbol o^n denotes the number of integral graphs of the corresponding order $o = 1, 2, \dots, 300$. In this table o^n is omitted if the corresponding number $n = 0$.

²The data given in Tables 1 and 2 are obtained in two different ways: (i) they are generated by using relations (2), (3) and (4); and (ii) by varying the parameters α, β, a, b in all possible ways in equation (1).

³In Tables 1 and 3 the number $\bar{\mu}_2$ denotes the second main eigenvalue of the corresponding integral graph $\overline{\alpha K_a \cup \beta K_{b,b}}$.

⁴For any integral graph $\overline{\alpha K_a \cup \beta K_{b,b}}$ with the largest eigenvalue $\bar{\mu}_1 = a + b$ we have (i) $\bar{\mu}_2 = -\frac{2\beta ab}{a+b}$ and (ii) $(a + b)(a + 2b + 1) = 2\beta b(2a + b)$ (see the proof of Theorem 1).

i	x_0	y_0	z	o	α	β	a	b	τ	t	k	ℓ	m	n	$\bar{\mu}_1$	$\bar{\mu}_2$
1	0	-1	1	10	1	1	8	1	4	1	2	1	1	1	4	-4
2	1	1	0	14	2	2	5	1	1	1	1	1	1	1	10	-3
3	1	0	1	16	10	1	1	3	1	1	1	1	3	1	14	-3
4	1	0	1	18	3	1	2	6	2	1	1	2	2	1	13	-4
5	-1	-1	1	20	2	1	9	1	3	3	1	1	1	1	12	-3
6	7	4	-1	20	2	1	7	3	1	1	1	1	3	1	14	-5
7	1	2	-1	22	1	1	18	2	2	1	5	1	2	1	9	-8
8	0	-1	1	22	1	3	16	1	8	2	2	1	1	2	12	-8
9	0	-1	2	22	2	3	8	1	4	1	2	1	1	1	16	-4
10	1	0	1	22	7	1	2	4	2	1	1	1	4	1	19	-4
11	0	-1	1	24	1	2	20	1	4	1	6	1	1	1	10	-8
12	0	-1	1	24	1	6	12	1	4	1	2	1	1	2	18	-8
13	-1	-1	1	26	3	2	6	2	2	1	1	1	2	1	21	-4
14	1	1	0	28	2	2	12	1	4	2	2	1	1	1	18	-4
15	-7	-3	-1	28	2	1	5	9	1	1	1	2	3	1	20	-7
16	-13	-2	-1	28	6	1	3	5	1	1	1	1	5	1	24	-5
17	3	8	-4	30	2	4	11	1	1	1	3	1	1	1	22	-5
18	1	1	0	30	1	5	10	2	2	1	1	1	2	3	25	-8
19	1	0	1	7	3	1	1	2	1	1	1	1	2	1	5	-2
20	1	1	0	9	1	1	5	2	1	1	1	1	2	1	5	-4
21	0	-1	1	10	1	2	6	1	2	1	2	1	1	1	6	-4
22	1	1	0	14	1	1	8	3	2	1	2	1	3	1	8	-6
23	1	0	1	14	6	1	1	4	1	1	2	1	4	1	11	-3
24	-1	-1	0	15	1	1	3	6	1	1	1	2	2	1	9	-4
25	1	0	2	15	7	2	1	2	1	1	1	1	2	1	13	-2
26	0	-1	1	16	1	3	10	1	2	1	4	1	1	1	10	-6
27	1	0	1	18	4	1	2	5	2	1	2	1	5	1	14	-4
28	3	4	-1	19	1	1	11	4	1	1	3	1	4	1	11	-8
29	0	-1	1	20	1	1	18	1	6	2	4	1	1	1	6	-6
30	1	3	-1	21	1	2	13	2	1	1	5	1	2	1	13	-8
31	1	0	1	21	9	1	1	6	1	1	3	1	6	1	17	-4
32	1	0	1	22	2	1	2	9	2	1	2	2	3	1	14	-4
33	0	-1	1	22	1	4	14	1	2	1	6	1	1	1	14	-8
34	0	-1	2	22	2	5	6	1	2	1	2	1	1	1	18	-4
35	1	0	3	23	11	3	1	2	1	1	1	1	2	1	21	-2
36	3	4	-1	24	1	1	14	5	2	1	4	1	5	1	14	-10
37	0	-1	1	26	1	1	20	3	10	3	2	2	1	1	12	-10
38	-3	-1	0	26	3	1	4	7	2	1	2	1	7	1	20	-6
39	3	5	-1	26	2	3	7	2	1	1	2	1	2	1	21	-5
40	0	-1	1	28	1	5	18	1	2	1	8	1	1	1	18	-10
41	1	0	1	28	12	1	1	8	1	1	4	1	8	1	23	-5
42	5	7	-2	29	1	1	17	6	1	1	5	1	6	1	17	-12
43	1	0	1	29	5	1	1	12	1	1	3	2	4	1	19	-4
44	1	0	1	29	3	1	3	10	3	2	1	3	2	1	21	-6
45	1	0	1	29	21	1	1	4	1	2	1	1	4	1	27	-4

TABLE 1

i	x_0	y_0	z	o	α	β	a	b	τ	t	k	ℓ	m	n	$\bar{\mu}_1$	$\bar{\mu}_2$
46	-1	-1	1	30	2	1	12	3	6	2	2	1	3	1	20	-6
47	1	0	2	30	14	2	1	4	1	1	2	1	4	1	27	-3
48	1	0	1	8	2	1	1	3	1	1	1	1	3	1	5	-2
49	0	-1	1	13	1	4	5	1	1	1	1	1	1	1	10	-4
50	-1	-1	0	14	1	1	2	6	1	1	1	2	3	1	8	-3
51	0	-1	1	16	1	2	12	1	3	2	2	1	1	1	8	-6
52	1	0	1	16	4	1	1	6	1	1	2	1	6	1	11	-3
53	0	-1	1	17	1	4	9	1	3	2	1	1	1	2	12	-6
54	1	0	2	17	5	2	1	3	1	1	1	1	3	1	14	-2
55	0	-1	1	18	1	1	14	2	7	4	1	2	1	1	8	-7
56	1	2	0	19	1	2	7	3	1	1	1	1	3	1	14	-6
57	0	-1	1	20	1	6	8	1	1	1	2	1	1	1	16	-6
58	-5	-2	-1	22	2	1	3	8	1	1	2	1	8	1	15	-5
59	1	0	1	22	12	1	1	5	1	2	1	1	5	1	19	-4
60	1	0	1	23	3	1	1	10	1	1	2	2	5	1	14	-3
61	1	0	1	24	6	1	1	9	1	1	3	1	9	1	17	-4
62	1	0	1	24	4	2	1	5	1	1	1	1	5	2	19	-2
63	1	0	3	26	8	3	1	3	1	1	1	1	3	1	23	-2
64	0	-1	1	27	1	2	15	3	5	3	1	3	1	1	18	-10
65	0	-1	1	27	1	8	11	1	1	1	3	1	1	1	22	-8
66	0	-1	1	28	1	4	20	1	5	3	2	1	1	2	16	-10
67	0	-1	2	28	2	9	5	1	1	1	1	1	1	1	25	-4
68	-3	-2	0	29	3	2	3	5	1	1	1	1	5	1	24	-4
69	0	-1	1	30	1	2	22	2	11	6	1	2	1	2	16	-11
70	-4	-1	-1	30	3	1	2	12	1	2	1	4	3	1	20	-5

TABLE 1. (continued)

007 ⁰¹	008 ⁰¹	009 ⁰¹	010 ⁰²	013 ⁰¹	014 ⁰⁴	015 ⁰²	016 ⁰⁴	017 ⁰²	018 ⁰³	019 ⁰²
020 ⁰⁴	021 ⁰²	022 ⁰⁹	023 ⁰²	024 ⁰⁵	026 ⁰⁵	027 ⁰²	028 ⁰⁷	029 ⁰⁵	030 ⁰⁶	031 ⁰⁵
032 ¹⁰	033 ⁰⁴	034 ²¹	035 ⁰⁴	036 ⁰⁷	037 ⁰²	038 ¹¹	039 ⁰²	040 ¹⁰	041 ⁰¹	042 ⁰⁶
043 ⁰⁷	044 ¹⁶	045 ⁰⁶	046 ²²	047 ⁰²	048 ¹²	049 ⁰⁵	050 ¹³	051 ⁰⁶	052 ¹⁴	053 ⁰⁴
054 ¹⁷	055 ⁰³	056 ¹⁰	057 ⁰⁵	058 ²²	059 ⁰⁶	060 ¹⁸	061 ¹⁰	062 ²⁷	063 ⁰⁶	064 ¹⁵
065 ⁰⁵	066 ¹⁹	067 ⁰⁷	068 ¹⁶	069 ⁰⁹	070 ¹⁸	071 ¹²	072 ¹²	073 ⁰⁸	074 ²⁹	075 ⁰³
076 ³⁴	077 ⁰⁴	078 ²⁴	079 ⁰⁷	080 ²⁰	081 ⁰⁴	082 ²⁹	083 ⁰⁶	084 ²²	085 ⁰⁴	086 ²³
087 ⁰⁶	088 ²²	089 ¹⁰	090 ²²	091 ⁰⁹	092 ²⁶	093 ¹⁴	094 ³⁴	095 ¹²	096 ³¹	097 ⁰⁹
098 ³³	099 ⁰⁹	100 ²¹	101 ⁰⁷	102 ³⁷	103 ¹³	104 ³⁰	105 ⁰⁷	106 ⁴⁶	107 ¹¹	108 ²⁹
109 ¹⁰	110 ²³	111 ¹¹	112 ²⁹	113 ⁰⁵	114 ³⁷	115 ⁰⁹	116 ³⁴	117 ⁰⁷	118 ⁴⁰	119 ¹¹

TABLE 2

120 ³⁰	121 ¹²	122 ³¹	123 ¹²	124 ⁴²	125 ¹¹	126 ³⁰	127 ¹¹	128 ³⁰	129 ¹⁵	130 ³⁷
131 ⁰⁸	132 ³⁶	133 ¹²	134 ⁴⁵	135 ¹²	136 ³⁹	137 ¹³	138 ⁴⁸	139 ¹⁵	140 ³⁵	141 ¹¹
142 ⁵⁰	143 ¹³	144 ³⁹	145 ¹⁰	146 ⁴²	147 ⁰⁷	148 ⁴⁴	149 ¹⁵	150 ³⁵	151 ⁰⁹	152 ³⁰
153 ¹⁶	154 ³³	155 ¹⁵	156 ⁴³	157 ¹⁴	158 ⁴⁷	159 ¹⁸	160 ⁴⁹	161 ¹⁰	162 ⁴⁷	163 ¹²
164 ⁵⁰	165 ¹⁰	166 ⁵⁷	167 ¹¹	168 ³⁹	169 ¹⁰	170 ³³	171 ⁰⁹	172 ⁵³	173 ¹⁰	174 ⁵⁰
175 ⁰⁸	176 ⁵¹	177 ⁰⁹	178 ⁵¹	179 ¹⁰	180 ³⁰	181 ¹²	182 ³⁵	183 ¹⁷	184 ⁴⁷	185 ⁰⁷
186 ⁴⁹	187 ¹²	188 ⁵⁶	189 ¹⁷	190 ⁶²	191 ¹⁷	192 ⁴⁰	193 ²¹	194 ⁶⁰	195 ¹⁹	196 ⁵³
197 ²⁰	198 ⁴⁷	199 ¹⁹	200 ³³	201 ¹³	202 ⁶¹	203 ¹⁴	204 ⁷⁶	205 ¹⁵	206 ⁵⁴	207 ¹⁸
208 ⁴⁹	209 ¹³	210 ⁴¹	211 ¹¹	212 ⁵⁸	213 ¹²	214 ⁶⁹	215 ¹⁵	216 ⁴⁷	217 ¹²	218 ⁵⁹
219 ¹⁴	220 ⁴⁹	221 ¹⁴	222 ⁶⁵	223 ¹³	224 ⁴⁰	225 ¹⁷	226 ⁶⁹	227 ¹⁸	228 ⁴⁸	229 ¹⁶
230 ⁵⁵	231 ²⁰	232 ⁴⁷	233 ¹⁸	234 ⁶⁰	235 ¹⁸	236 ⁵⁵	237 ²⁰	238 ⁶⁴	239 ¹³	240 ⁵⁵
241 ²⁴	242 ⁶⁴	243 ¹³	244 ⁷⁴	245 ¹³	246 ⁶⁸	247 ¹¹	248 ⁵⁶	249 ²⁵	250 ⁷³	251 ¹⁶
252 ⁵³	253 ²⁰	254 ⁶⁸	255 ²²	256 ⁵⁷	257 ¹⁰	258 ⁷³	259 ¹⁶	260 ⁵⁷	261 ²²	262 ⁵⁵
263 ¹⁷	264 ⁶⁶	265 ¹⁶	266 ⁵⁰	267 ¹²	268 ⁶⁶	269 ¹⁴	270 ⁵¹	271 ¹⁷	272 ⁵⁷	273 ²¹
274 ⁷¹	275 ¹⁹	276 ⁸³	277 ¹⁷	278 ⁶⁵	279 ²⁸	280 ⁵²	281 ¹⁷	282 ⁷⁵	283 ²⁰	284 ⁸⁴
285 ²⁴	286 ⁷²	287 ¹⁹	288 ⁶⁰	289 ¹⁵	290 ⁶²	291 ²³	292 ⁶⁶	293 ¹⁰	294 ⁷⁷	295 ¹⁶
296 ⁸⁰	297 ¹³	298 ⁷⁰	299 ¹⁴	300 ⁶⁷						

TABLE 2. (continued)

a	b	o	$\bar{\mu}_1$	$\bar{\mu}_2$
7865	585	12545	8450	-4356
53492	5676	87548	59168	-30789
7024874	745390	11497214	7770264	-4043315
127230675	13500094	208231239	140730769	-73230300
480286984490	35719102710	766039806170	516006087200	-265972368231
12529086263859	931792310790	19983424750179	13460878574649	-6938332399120

TABLE 3

Table 3 contains the integral graphs $\overline{\alpha K_a \cup \beta K_{b,b}}$ with $\bar{\mu}_1 = a + b$ and $a > (b + 1)$, obtained from the classes represented in Propositions 2 and 3 for $i = 0$. We note that any graph in this list is an integral graph with the minimal number of vertices for the corresponding class. The first, second, . . . , sixth integral graph in List 3 belongs to the class described in Proposition m (n^0), where $(m, n) = (3, 2), (2, 2), (2, 3), (2, 1), (3, 3)$ and $(3, 1)$, respectively.

Acknowledgement. The author is very grateful to prof. Dragoš Cvetković for his valuable comments and suggestions concerning this paper.

References

- [1] D. Cvetković, M. Doob, H. Sachs, *Spectra of graphs – Theory and applications*, 3rd revised and enlarged edition, J. A. Barth Verlag, Heidelberg – Leipzig, 1995.
- [2] M. Lepović, *On integral graphs which belong to the class $\overline{\alpha K_a \cup \beta K_b}$* , submitted to Linear and Multilinear Algebra.
- [3] M. Lepović, *On integral graphs which belong to the class $\overline{\alpha K_{a,b}}$* , Graphs and Combinatorics, in press.
- [4] M. Lepović, *On integral graphs which belong to the class $\overline{\alpha K_{a,a} \cup \beta K_{b,b}}$* , submitted to Indian J. Pure Appl. Math.
- [5] M. Lepović, *On integral graphs which belong to the class $\overline{\alpha K_a \cup \beta K_{b,b}}$* , submitted to Discrete Math.
- [6] D. Redmond, *Number Theory – an Introduction*, Marcel Dekker, New York, 1996.

Prirodno-matematički fakultet
34000 Kragujevac
Serbia and Montenegro
lepovic@knez.uis.kg.ac.yu

(Received 12 11 2002)
(Revised 24 12 2003)