

## CHARACTERIZATIONS OF $sn$ -METRIZABLE SPACES

Ying Ge

*Communicated by Rade Živaljević*

ABSTRACT. We give some characterizations of  $sn$ -metrizable spaces. We prove that a space is an  $sn$ -metrizable space if and only if it has a locally-finite point-star  $sn$ -network. As an application of the result, a space is an  $sn$ -metrizable space if and only if it is a sequentially quotient,  $\pi$  (compact),  $\sigma$ -image of a metric space.

### 1. Introduction

To characterize generalized metric spaces by some point-star networks and by some map images of metric spaces are important questions in general topology. The former was due to thought of Alexandroff and Urysohn characterized metric spaces by point-star neighborhood base [1], and the later came from the Alexandroff's suggestion to investigate images or preimages of “nice” spaces under “nice” maps [2]. Having gained some enlightenment from investigation of  $g$ -developable spaces by Tanaka in [18], Lin characterized  $g$ -metrizable spaces by point-star weak neighborhood base, and proved that a space  $X$  is  $g$ -metrizable if and only if  $X$  has a locally-finite point-star weak neighborhood base [9]. Notice that it is difficult to characterize generalized metric spaces by using general map images of metric spaces. Recently the authors of [13] introduced  $\sigma$ -map, and proved that a space  $X$  is  $g$ -metrizable if and only if  $X$  is a compact-covering, quotient, compact,  $\sigma$ -image of a metric space [13]. In a private communication the first author of [13] posed the question if there is a similar characterization for  $sn$ -metrizable spaces, an important generalization of  $g$ -metrizable spaces. In this paper, we investigate this question, prove that a space is an  $sn$ -metrizable space if and only if it has a locally-finite point-star  $sn$ -network, if and only if it is a sequentially quotient,  $\pi$  (compact),  $\sigma$ -image of a metric space. As a corollary of the results above, a space

---

2000 *Mathematics Subject Classification*: 54C10, 54D50, 54D55, 54E99.

*Key words and phrases*:  $sn$ -metrizable space, point star network,  $sn$ -network,  $\sigma$ -map, sequentially quotient map,  $\pi$ -map.

The author was supported in part by NSF of the Education Committee of Jiangsu Province in China No. 02KJB110001.

is a  $g$ -metrizable space if and only if it is a quotient,  $\pi$  (compact),  $\sigma$ -image of a metric space, which improves a foregoing result on  $g$ -metrizable spaces by omitting the condition “compact-covering” in the statement.

Throughout this paper, all spaces are assumed to be regular  $T_1$ , and all maps are continuous and onto. Every convergent sequence contains its limit point.  $N$  denotes the set of all natural numbers.  $S_1$  denotes the subspace  $\{0\} \cup \{1/n : n \in N\}$  of the real numbers space. Let  $A$  be a subset of a space  $X$ .  $\bar{A}$  denotes the closure of  $A$ . Furthermore let  $x \in X$ ,  $\mathcal{U}$  be a collection of subsets of  $X$ ,  $\bigcup \mathcal{U} = \bigcup \{U : U \in \mathcal{U}\}$ ,  $\text{st}(x, \mathcal{U}) = \bigcup \{U \in \mathcal{U} : x \in U\}$ . The sequence  $\{x_n : n \in N\}$ , the sequence  $\{P_n : n \in N\}$  of subsets and the sequence  $\{\mathcal{P}_n : n \in N\}$  of collections of subsets are abbreviated to  $\{x_n\}$ ,  $\{P_n\}$  and  $\{\mathcal{P}_n\}$  respectively. For terms which are not defined here, please refer to [9].

DEFINITION 1.1. [5] Let  $X$  be a space,  $x \in P \subset X$ .  $P$  is a sequential neighborhood of  $x$ , if every sequence  $\{x_n\}$  converging to  $x$  is eventually in  $P$ , i.e., there is  $k \in N$  such that  $x_n \in P$  for  $n > k$ .

REMARK 1.2. (1)  $P$  is a sequential neighborhood of  $x$  if and only if  $x \in P$  and every sequence  $\{x_n\}$  converging to  $x$  is frequently in  $P$ , i.e., for every  $k \in N$ , there is  $n > k$  such that  $x_n \in P$ .

(2) The intersection of finitely many sequential neighborhoods of  $x$  is a sequential neighborhood of  $x$

DEFINITION 1.3. [6], [10] A cover  $\mathcal{P} = \bigcup \{\mathcal{P}_x : x \in X\}$  of a space  $X$  is a  $cs$ -network ( $cs^*$ -network), if every convergent sequence  $S$  converging to a point  $x \in U$  with  $U$  open in  $X$ , then  $S$  is eventually (frequently) in  $P \subset U$  for some  $P \in \mathcal{P}_x$ . A space  $X$  is  $csf$ -countable, if  $X$  has a  $cs$ -network  $\mathcal{P} = \bigcup \{\mathcal{P}_x : x \in X\}$  such that  $\mathcal{P}_x$  is countable for every  $x \in X$ ; is an  $\aleph$ -space, if  $X$  has a  $\sigma$ -locally-finite  $cs$ -network.

DEFINITION 1.4. [3], [11] Let  $\mathcal{P} = \bigcup \{\mathcal{P}_x : x \in X\}$  be a cover of a space  $X$ . Assume that  $\mathcal{P}$  satisfies the following (a) and (b) for every  $x \in X$ .

(a)  $\mathcal{P}$  is a network of  $X$ , that is, whenever  $x \in U$  with  $U$  open in  $X$ , then  $x \in P \subset U$  for some  $P \in \mathcal{P}_x$ ;  $\mathcal{P}_x$  is said to be a network at  $x$  for every  $x \in X$ .

(b) If  $P_1, P_2 \in \mathcal{P}_x$ , then there exists  $P \in \mathcal{P}_x$  such that  $P \subset P_1 \cap P_2$ .

(1)  $\mathcal{P}$  is a weak base of  $X$ , if for  $G \subset X$ ,  $G$  is open in  $X$  if and only if for every  $x \in G$  there exists  $P \in \mathcal{P}_x$  such that  $P \subset G$ ;  $\mathcal{P}_x$  is said to be a weak neighborhood base ( $wn$ -base for short) at  $x$ .

(2)  $\mathcal{P}$  is an  $sn$ -network of  $X$ , if every element of  $\mathcal{P}_x$  is a sequential neighborhood of  $x$  for every  $x \in X$ ;  $\mathcal{P}_x$  is said to be an  $sn$ -network at  $x$ .

(3) A space  $X$  is  $gf$ -countable ( $snf$ -countable), if  $X$  has a weak base ( $sn$ -network)  $\mathcal{P} = \bigcup \{\mathcal{P}_x : x \in X\}$  such that  $\mathcal{P}_x$  is countable for every  $x \in X$ ; is  $g$ -metrizable ( $sn$ -metrizable), if  $X$  has a  $\sigma$ -locally-finite weak base ( $sn$ -network).

REMARK 1.5. [8], [10], [11] (1) weak base  $\Rightarrow sn$ -network  $\Rightarrow cs$ -network, so  $gf$ -countable  $\Rightarrow snf$ -countable  $\Rightarrow csf$ -countable, and  $g$ -metrizable  $\Rightarrow sn$ -metrizable  $\Rightarrow \aleph$ .

(2)  $g$ -metrizable  $\Leftrightarrow k$  and  $sn$ -metrizable.

(3) In the class of Fréchet spaces, metrizable  $\Leftrightarrow g$ -metrizable  $\Leftrightarrow sn$ -metrizable.

DEFINITION 1.6. [12] Let  $\{\mathcal{P}_n\}$  be a sequence of covers of a space  $X$ .

- (1)  $\{\mathcal{P}_n\}$  is a point-star network (point-star  $sn$ -network, point-star  $wn$ -base) of  $X$ , if  $\{\text{st}(x, \mathcal{P}_n)\}$  is a network ( $sn$ -network,  $wn$ -base) at  $x$  for every  $x \in X$ .
- (2)  $\{\mathcal{P}_n\}$  is locally-finite (hereditarily closure-preserving, discrete, point-finite), if every  $\mathcal{P}_n$  is locally-finite (hereditarily closure-preserving, discrete, point-finite).

DEFINITION 1.7. [4], [13], [14], [17] Let  $f : X \rightarrow Y$  be a map.

- (1) Let  $(X, d)$  be a metric space.  $f$  is a  $\pi$ -map, if  $d(f^{-1}(y), X - f^{-1}(U)) > 0$  for every  $y \in Y$  and every neighborhood  $U$  of  $y$  in  $Y$ .
- (2)  $f$  is a  $\sigma$ -map, if there exists a base  $\mathcal{B}$  of  $X$  such that  $f(\mathcal{B})$  is  $\sigma$ -locally-finite in  $Y$ .
- (3)  $f$  is a sequentially-quotient map, if whenever  $S$  is a convergent sequence in  $Y$  there is a convergent sequence  $L$  in  $X$  such that  $f(L)$  is a subsequence of  $S$ .
- (4)  $f$  is a quotient map, if whenever  $U \subset Y$ ,  $f^{-1}(U)$  is open in  $X$  iff  $U$  is open in  $Y$ .

REMARK 1.8. [9] (1) Closed map  $\Rightarrow$  quotient map.

- (2) If the domain is sequential, then quotient map  $\Rightarrow$  sequentially-quotient map.
- (3) If the image is sequential, then sequentially-quotient map  $\Rightarrow$  quotient map.

## 2. $sn$ -metrizable spaces and point-star $sn$ -networks

Let  $\mathcal{P}$  be a collection of subsets of a space  $X$ . We recall that  $\mathcal{P}$  is a  $k$ -network of  $X$ , if whenever  $K$  is a compact subset of an open set  $U$ , there is a finite  $\mathcal{F} \subset \mathcal{P}$  such that  $K \subset \bigcup \mathcal{F} \subset U$ .

LEMMA 2.1. [9] Let  $\mathcal{P}$  be a  $\sigma$ -hereditarily closure-preserving collection of subsets of a space  $X$ . If  $\mathcal{P}$  is a  $cs^*$ -network of  $X$ , then  $\mathcal{P}$  is a  $k$ -network of  $X$ .

LEMMA 2.2. [7], [8] For a space  $X$ , the following are equivalent.

- (1)  $X$  is an  $sn$ -metrizable space.
- (2)  $X$  has a  $\sigma$ -discrete (closed)  $sn$ -network.
- (3)  $X$  has a  $\sigma$ -hereditarily closure-preserving (closed)  $sn$ -network.
- (4)  $X$  is an  $snf$ -countable space with a  $\sigma$ -hereditarily closure-preserving (closed)  $k$ -network.

The following lemma can be obtained from Lemma 3.1 and Lemma 3.2.

LEMMA 2.3. A space is  $sn$ -metrizable if and only if it is an  $snf$ -countable space with a  $\sigma$ -hereditarily closure-preserving (closed)  $cs^*$ -network.

LEMMA 2.4. Let  $\mathcal{P}$  be a hereditarily closure-preserving collection of subsets of a space  $X$ , and  $L$  be a convergent sequence, which is eventually in  $\bigcup \mathcal{P}$ . Then there exists  $P \in \mathcal{P}$  such that  $L$  is frequently in  $P$ .

PROOF. If not, then  $L \cap P$  is finite for every  $P \in \mathcal{P}$ . We may assume without loss of generality that  $L = \{x_n : n \in \mathbb{N}\} \subset \bigcup \mathcal{P}$  and  $L$  is infinite. Pick  $x_{n_1} \in \bigcup \mathcal{P}$ , then there exists  $P_1 \in \mathcal{P}$  such that  $x_{n_1} \in P_1$ . Since  $L$  is infinite and  $L \cap P_1$  is

finite,  $L - P_1$  is infinite, we may pick  $n_2 > n_1$  and  $P_2 \in \mathcal{P}$  such that  $x_{n_2} \in P_2$  and  $x_{n_2} \neq x_{n_1}$ . By induction, we can obtain a convergent subsequence  $\{x_{n_k}\}$  of  $L$  such that  $x_{n_k} \in P_k \in \mathcal{P}$  for every  $k \in N$ , and  $x_{n_k} \neq x_{n_l}$  if  $k \neq l$ . Thus  $\{\{x_{n_k}\} : k \in N\}$  is not closure-preserving. This contradicts the fact that  $\mathcal{P}$  is hereditarily closure-preserving.  $\square$

**THEOREM 2.5.** *For a space  $X$ , the following are equivalent.*

- (1)  $X$  is an  $sn$ -metrizable space.
- (2)  $X$  has a locally-finite point-star  $sn$ -network.
- (3)  $X$  has a hereditarily closure-preserving point-star  $sn$ -network.

**PROOF.** (1)  $\Rightarrow$  (2): Let  $X$  is an  $sn$ -metrizable space.  $X$  has an  $sn$ -network  $\mathcal{P} = \bigcup\{\mathcal{P}_n : n \in N\}$  from Lemma 2.2, where every  $\mathcal{P}_n$  is a discrete collection of closed subsets of  $X$ . Write  $\mathcal{P} = \bigcup\{\mathcal{B}_x : x \in X\}$ , where every  $\mathcal{B}_x$  is an  $sn$ -network at  $x$ . For every  $n \in N$ , Put  $F_n = \{x \in X : \mathcal{P}_n \cap \mathcal{B}_x = \emptyset\}$ ,  $\mathcal{F}_n = \{F_n\} \cup \mathcal{P}_n$ . Obviously,  $\{\mathcal{F}_n\}$  is a sequence of locally-finite covers of  $X$ . Let  $x \in X$ . We prove  $\mathcal{F}_x = \{\text{st}(x, \mathcal{F}_n) : n \in N\}$  is an  $sn$ -network at  $x$  as follows.

**Claim A.**  $\mathcal{F}_x$  is a network at  $x$ : Let  $x \in X$  and  $U$  be an open neighborhood of  $x$ . Since  $\mathcal{B}_x$  is a network at  $x$ , there exists  $P_x \in \mathcal{B}_x \cap \mathcal{P}_n$  for some  $n \in N$ . such that  $x \in P_x \in U$ . Then  $x \notin F_n$ . Notice that elements of  $\mathcal{P}_n$  are mutually disjoint. So  $P_x = \text{st}(x, \mathcal{F}_n)$ , that is,  $x \in \text{st}(x, \mathcal{F}_n) \subset U$ . This proves that  $\mathcal{F}_x$  is a network at  $x$ .

**Claim B.**  $\mathcal{F}_x$  satisfies Definition 1.4(b): Consider  $\text{st}(x, \mathcal{F}_n)$  and  $\text{st}(x, \mathcal{F}_m)$ , where  $n, m \in N$ . Then one of the following holds:

- (a)  $\mathcal{B}_x \cap \mathcal{P}_n \neq \emptyset$ ,  $\mathcal{B}_x \cap \mathcal{P}_m \neq \emptyset$ ;
- (b)  $\mathcal{B}_x \cap \mathcal{P}_n \neq \emptyset$ ,  $\mathcal{B}_x \cap \mathcal{P}_m = \emptyset$ ;
- (c)  $\mathcal{B}_x \cap \mathcal{P}_n = \emptyset$ ,  $\mathcal{B}_x \cap \mathcal{P}_m \neq \emptyset$ ;
- (d)  $\mathcal{B}_x \cap \mathcal{P}_n = \emptyset$ ,  $\mathcal{B}_x \cap \mathcal{P}_m = \emptyset$ .

We prove the case (b), the other cases are similar.  $\mathcal{B}_x \cap \mathcal{P}_n \neq \emptyset$ , so there exists  $P_n \in \mathcal{B}_x \cap \mathcal{P}_n$ . Notice that  $\mathcal{B}_x \cap \mathcal{P}_m = \emptyset$ .  $U = X - \bigcup\{P \in \mathcal{P}_m : x \notin P\}$  is an open neighborhood of  $x$ , and  $U \subset \text{st}(x, \mathcal{F}_m)$ , so there exists  $P_l \in \mathcal{B}_x \cap \mathcal{P}_l$  such that  $P_l \subset U \subset \text{st}(x, \mathcal{F}_m)$ . There exists  $P_k \in \mathcal{B}_x \cap \mathcal{P}_k$  such that  $P_k \subset P_n \cap P_l$ .  $\mathcal{B}_x \cap \mathcal{P}_n \neq \emptyset$  and  $\mathcal{B}_x \cap \mathcal{P}_k \neq \emptyset$ , so  $x \notin F_n$  and  $x \notin F_k$ . Notice that  $\mathcal{P}_n$  and  $\mathcal{P}_k$  are discrete.  $P_n = \text{st}(x, \mathcal{F}_n)$  and  $P_k = \text{st}(x, \mathcal{F}_k)$ . This proves that  $\text{st}(x, \mathcal{F}_k) = P_k \subset P_n \cap P_l \subset \text{st}(x, \mathcal{F}_n) \cap \text{st}(x, \mathcal{F}_m)$ .

**Claim C.**  $\text{st}(x, \mathcal{F}_n)$  is a sequential neighborhood of  $x$  for every  $n \in N$ : Let  $L$  is a sequence converging to  $x \in \text{st}(x, \mathcal{F}_n)$ . If  $\mathcal{B}_x \cap \mathcal{P}_n \neq \emptyset$ , then there exists  $P \in \mathcal{B}_x \cap \mathcal{P}_n$ . Since  $P$  is a sequential neighborhood of  $x$ ,  $L$  is eventually in  $P$ , thus  $L$  is eventually in  $\text{st}(x, \mathcal{F}_n)$ . If  $\mathcal{B}_x \cap \mathcal{P}_n = \emptyset$ , Put  $U = X - \bigcup\{P \in \mathcal{P}_n : x \notin P\}$ . Then  $U$  is an open neighborhood of  $x$ , so  $L$  is eventually in  $U$ . It is easy to see that  $U \subset \text{st}(x, \mathcal{F}_n)$ , So  $L$  is eventually in  $\text{st}(x, \mathcal{F}_n)$ . This proves that  $\text{st}(x, \mathcal{F}_n)$  is a sequential neighborhood of  $x$  for every  $n \in N$ .

By the three claims above,  $\{\text{st}(x, \mathcal{F}_n) : n \in N\}$  is an  $sn$ -network at  $x$ . This proves that  $\{\mathcal{F}_n\}$  is a locally-finite point-star  $sn$ -network.

(2)  $\Rightarrow$  (3): This is clear.

(3)  $\Rightarrow$  (1): Let  $X$  has a hereditarily closure-preserving point-star  $sn$ -network  $\{\mathcal{F}_n\}$ . Then  $\mathcal{F}_n$  is hereditarily closure-preserving for every  $n \in N$ . It is easy to see that  $X$  is  $snf$ -countable. We only need prove that  $\bigcup\{\mathcal{F}_n : n \in N\}$  is a  $cs^*$ -network of  $X$  by Lemma 2.3. Let  $L$  be a sequence in  $X$  converging to  $x$ , and  $U$  be an open

neighborhood of  $x$ .  $\{st(x, \mathcal{F}_n) : n \in N\}$  is a network at  $x$ , so there exists  $n \in N$  such that  $x \in st(x, \mathcal{F}_n) \subset U$ . Since  $st(x, \mathcal{F}_n)$  is a sequential neighborhood of  $x$ ,  $L$  is eventually in  $st(x, \mathcal{F}_n)$ . By Lemma 2.4, there exists  $F \in \mathcal{F}_n$  and  $x \in F$  such that  $L$  frequently in  $F$ . So there exists a subsequence  $S$  of  $L$  such that  $S \subset F$ . This proves that  $\bigcup\{\mathcal{F}_n : n \in N\}$  is a  $cs^*$ -network of  $X$ .  $\square$

Can “locally-finite” in Theorem 2.5 be replaced by “discrete” or “point-finite”? The answers are negative by the following.

EXAMPLE 2.6. There exists a space  $X$  with a point-finite point-star  $wn$ -base, and  $X$  is not an  $\aleph$ -space.

PROOF. Let  $I$  denote the closed interval  $[0, 1]$ , let  $S(x)$  be homeomorphic to  $S_1$  for every  $x \in I$ , and  $T = \bigoplus\{S(x) : x \in I\}$ .  $X$  is the quotient space obtained from the topological sum  $Z = I \oplus T$  by identifying every  $x \in I$  with the limit point of  $S(x)$ . Then  $Z$  is a locally compact metric space, and the natural map  $f : Z \rightarrow X$  is quotient, compact, compact-covering. So  $X$  has a point-finite point-star  $wn$ -base by [9, Theorem 2.9.14] and  $X$  has not point-countable  $cs$ -networks by [9, Example 2.9.27].  $\square$

LEMMA 2.7. [16] *A space is metrizable if and only if it has a locally-finite point-star closed network.*

PROPOSITION 2.8. *Let  $X$  have a discrete point-star network. Then  $X$  is metrizable.*

PROOF. Let  $\{\mathcal{P}_n\}$  is a point-star network of  $X$ , where every  $\mathcal{P}_n$  is discrete. Put  $\overline{\mathcal{P}}_n = \{\overline{P} : P \in \mathcal{P}_n\}$ ; then  $\{\overline{\mathcal{P}}_n\}$  is discrete point-star closed network of  $X$ . Thus  $X$  is metrizable by Lemma 2.7.  $\square$

### 3. $sn$ -metrizable spaces and images of metric spaces

LEMMA 3.1. *Let  $X$  be a sequentially-quotient,  $\pi$ -image of a metric space. Then  $X$  has a point-star  $sn$ -network, and so  $X$  is  $snf$ -countable.*

PROOF. Let  $f : M \rightarrow X$  be a sequentially-quotient,  $\pi$ -map,  $(M, d)$  be a metric space. Put  $\mathcal{P}_n = \{f(B(z, 1/n)) : z \in M\}$  for every  $n \in N$ .

(a)  $\{\mathcal{P}_n\}$  is a point-star network of  $X$ : Let  $x \in U$  with  $U$  open in  $X$ . Since  $f$  is a  $\pi$ -map, there exists  $n \in N$  such that  $d(f^{-1}(x), M - f^{-1}(U)) > 1/n$ . Pick  $m \in N$  such that  $m \geq 2n$ . Notice that  $\mathcal{P}_m$  covers  $X$ , there exists  $z \in M$  such that  $x \in f(B(z, 1/m))$ . It is easy to see that  $f^{-1}(x) \cap B(z, 1/m) \neq \emptyset$ . We claim that  $B(z, 1/m) \subset f^{-1}(U)$ . If not, there would exist  $y \in B(z, 1/m) \cap (M - f^{-1}(U))$ . Pick  $t \in f^{-1}(x) \cap B(z, 1/m)$ , then  $d(t, y) \leq d(t, z) + d(z, y) < 2/m \leq 1/n$ . This contradicts the fact that  $d(f^{-1}(x), M - f^{-1}(U)) > 1/n$ , hence  $B(z, 1/m) \subset f^{-1}(U)$ . Thus  $st(x, \mathcal{P}_m) \subset U$ , so  $\{\mathcal{P}_n\}$  is a point-star network of  $X$ .

(b)  $st(x, \mathcal{P}_n)$  is a sequential neighborhood of  $x$  for every  $x \in X$  and every  $n \in N$ : Let  $S$  be a sequence converging to  $x$ . Since  $f$  is sequentially-quotient, there exists a sequence  $L$  converging to  $t \in f^{-1}(x) \subset M$  such that  $f(L) = S'$  is a subsequence

of  $S$ . Write  $B = B(t, 1/n)$ , then  $f(B) \in \mathcal{P}_n$ . Since  $B$  is an open neighborhood of  $t$ ,  $L$  is eventually in  $B$ , and so  $S' = f(L)$  is eventually in  $f(B) \subset \text{st}(x, \mathcal{P}_n)$ . That is,  $S$  is frequently in  $\text{st}(x, \mathcal{P}_n)$ , so  $\text{st}(x, \mathcal{P}_n)$  is a sequential neighborhood of  $x$ .

By the above,  $\{\mathcal{P}_n\}$  is a point-star  $sn$ -network of  $X$ .  $\square$

As an application of the lemma above we have the following proposition, which improves a typical result that perfect maps preserve metric spaces.

PROPOSITION 3.2. *Closed,  $\pi$ -maps preserve metric spaces.*

PROOF. Let  $f : X \rightarrow Y$  be a closed,  $\pi$ -map, and  $X$  be a metric space. Then  $Y$  is a Fréchet space with  $\sigma$ -hereditarily closure-preserving  $k$ -network. Since closed maps are sequentially-quotient maps (Remark 1.8),  $Y$  is  $snf$ -countable from Lemma 3.1, hence  $Y$  is  $sn$ -metrizable by Lemma 2.2. So  $Y$  is a metric space by Remark 1.5.  $\square$

LEMMA 3.3. *Let  $f : X \rightarrow Y$  be a map,  $\{y_n\}$  a sequence converging to  $y \in Y$ . If  $\{B_k\}$  is a decreasing network at some point  $x \in f^{-1}(y)$ , and  $\{y_n\}$  is frequently in  $f(B_k)$  for every  $k \in N$ , then there exists a sequence  $\{x_k\}$  converging to  $x$  such that  $\{f(x_k)\}$  is a subsequence of  $\{y_n\}$ .*

PROOF. Since  $\{y_n\}$  is frequently in  $f(B_1)$ , there exists  $n_1 \in N$  such that  $y_{n_1} \in f(B_1)$ . Pick  $x_1 \in f^{-1}(y_{n_1}) \cap B_1$ . We construct a sequence  $\{x_k\}$  by induction as follows. Assume  $\{x_k\}$  have been picked for  $k \in N$ . Since  $\{y_n\}$  is frequently in  $f(B_{k+1})$ , there exists  $n_{k+1} \in N$  and  $n_{k+1} > n_k$  such that  $y_{n_{k+1}} \in f(B_{k+1})$ , and so we may pick  $x_{k+1} \in f^{-1}(y_{n_{k+1}}) \cap B_{k+1}$ . Thus we construct a sequence  $\{x_k\}$ . It is easy to see that  $\{f(x_k)\} = \{y_{n_k}\}$  is a subsequence of  $\{y_n\}$ . Notice that  $x_k \in B_k$  for every  $k \in N$ , and  $\{B_k\}$  is a decreasing network at  $x$ . So  $\{x_k\}$  converges to  $x$ .  $\square$

THEOREM 3.4. *For a space  $X$ , the following are equivalent.*

- (1)  $X$  is an  $sn$ -metrizable space.
- (2)  $X$  is a sequentially-quotient, compact,  $\sigma$ -image of a metric space.
- (3)  $X$  is a sequentially-quotient,  $\pi$ ,  $\sigma$ -image of a metric space.

PROOF. We only need prove (1)  $\Rightarrow$  (2) and (3)  $\Rightarrow$  (1).

(1)  $\Rightarrow$  (2): Since  $X$  is an  $sn$ -metrizable space,  $X$  has a locally-finite point-star  $sn$ -network  $\{\mathcal{P}_n\}$  by Theorem 2.5. Write  $\mathcal{P}_n = \{P_\alpha : \alpha \in A_n\}$  for every  $n \in N$ , where  $\{A_n\}$  are mutually disjoint. For every  $n \in N$ , put  $\mathcal{F}_n = \{\bigcap_{i \leq n} P_{\alpha_i} : \alpha_i \in A_i, i = 1, 2, \dots, n\}$ ; then  $\mathcal{F}_n$  is locally-finite. Endow the discrete topology on  $A_n$  for every  $n \in N$ . Put  $Z = \{a = (\alpha_n) \in \prod_{n \in N} A_n : \{P_{\alpha_n}\} \text{ is a network at an } x_a \in X\}$ . Then  $Z$  is a metric space, where metric  $d$  is as follows: Let  $a = (\alpha_n), b = (\beta_n) \in Z$ ,  $d(a, b) = 0$  if  $a = b$ , and  $d(a, b) = 1/\min\{n \in N : \alpha_n \neq \beta_n\}$  if  $a \neq b$ .

Obviously,  $\{P_{\alpha_n}\}$  is a network at some  $x_a \in X$  iff  $x_a \in \bigcap_{n \in N} P_{\alpha_n}$  for  $a = (\alpha_n) \in \prod_{n \in N} A_n$ . It is not difficulty to prove that  $f : Z \rightarrow X$  defined by  $f(a) = x_a$  is a map. We only need prove that  $f$  is a sequentially-quotient, compact,  $\sigma$ -map.

(a)  $f$  is a sequentially-quotient map: Let  $x \in X$ . Assume  $S$  is a sequence converging to  $x$ . For every  $n \in N$ , since  $\text{st}(x, \mathcal{P}_n)$  is a sequential neighborhood of  $x$ ,  $S$  is eventually in  $\text{st}(x, \mathcal{P}_n)$ . Notice that  $\mathcal{P}_n$  is point-finite, so there exists a subsequence  $S'$  of  $S$  such that  $S'$  is eventually in some element of  $\mathcal{P}_n$ .

Let  $L = L_0 = \{x_n : n \in N\} \cup \{x\}$  be a sequence converging to  $x$ . By induction, for every  $n \in N$ , we may choose  $\alpha_n \in A_n$  and a subsequence  $L_n$  of  $L_0$ , such that  $L_n$  is a subsequence of  $L_{n-1}$ , and  $L_n$  is eventually in  $P_{\alpha_n} \in \mathcal{P}_n$ . Put  $z = (\alpha_n) \in \prod_{n \in N} A_n$ . Obviously,  $\{P_{\alpha_n} : n \in N\}$  is a network at  $x$ , so  $z \in Z$  and  $f(z) = x$ . Put  $Z_n = \{(\beta_k) \in Z : \beta_k = \alpha_k \text{ for } k \leq n\}$ . Then  $\{Z_n\}$  is a decreasing neighborhood base at  $z$ . We prove that  $f(Z_n) = \bigcap_{k \leq n} P_{\alpha_k}$  for every  $n \in N$  as follows.

In fact, let  $b = (\beta_k) \in Z_n$ . Then  $f(b) \in \bigcap_{k \in N} P_{\beta_k} \subset \bigcap_{k \leq n} P_{\alpha_k}$ , so  $f(Z_n) \subset \bigcap_{k \leq n} P_{\alpha_k}$ . On the other hand, let  $y \in \bigcap_{k \leq n} P_{\alpha_k}$ . Then there exists  $c' = (\gamma'_k) \in Z$  such that  $f(c') = y$ . For every  $k \in N$ , put  $\gamma_k = \alpha_k$  if  $k \leq n$ , and  $\gamma_k = \gamma'_k$  if  $k > n$ . Put  $c = (\gamma_k)$ . It is easy to see that  $y \in \bigcap_{n \in N} P_{\gamma_n}$ , so  $c \in Z$  and  $f(c) = y$ , hence  $c \in Z_n$ . This shows that  $y \in f(Z_n)$ . So  $\bigcap_{k \leq n} P_{\alpha_k} \subset f(Z_n)$ . Thus  $f(Z_n) = \bigcap_{k \leq n} P_{\alpha_k}$ .

Let  $n \in N$ . By the construction of  $L_n$ ,  $L_n$  is eventually in  $P_{\alpha_k}$  for every  $k \leq n$ , and so  $L_n$  is eventually in  $\bigcap_{k \leq n} P_{\alpha_k} = f(Z_n)$ . Thus  $L$  is frequently in  $f(Z_n)$  for every  $n \in N$ . By Lemma 3.3, there exists a sequence  $\{z_n\}$  converging to  $z$ , and  $\{f(z_n)\}$  is a subsequence of  $L$ . So  $f$  is sequentially-quotient map.

(b)  $f$  is a compact map: Let  $x \in X$ . Put  $B_n = \{\alpha \in A_n : x \in P_\alpha\}$ , then  $\prod_{n \in N} B_n$  is a compact subset of  $\prod_{n \in N} A_n$ . It is easy to prove that  $f^{-1}(x) = \prod_{n \in N} B_n$ . So  $f$  is a compact map.

(c)  $f$  is a  $\sigma$ -map: Put  $B(\alpha_1, \alpha_2, \dots, \alpha_n) = \{(\beta_k) \in Z : \beta_k = \alpha_k \text{ for every } k \leq n\}$ , where  $(\alpha_i) \in Z$  and  $n \in N$ . Then  $\{B(\alpha_1, \alpha_2, \dots, \alpha_n) : (\alpha_i) \in Z \text{ and } n \in N\}$  is a base of  $Z$ . By an argument similar to that in (a),  $f(B(\alpha_1, \alpha_2, \dots, \alpha_n)) = \bigcap_{k \leq n} P_{\alpha_k} \in \mathcal{F}_n$ . Notice that  $\mathcal{F}_n$  is locally-finite. So  $\{f(B(\alpha_1, \alpha_2, \dots, \alpha_n)) : (\alpha_i) \in Z \text{ and } n \in N\}$  is  $\sigma$ -locally-finite. Thus  $f$  is a  $\sigma$ -map.

(3)  $\Rightarrow$  (1): Let  $Z$  be a metric space, and  $f : Z \rightarrow X$  a sequentially-quotient,  $\pi$ ,  $\sigma$ -map. Then  $X$  is  $snf$ -countable by Lemma 3.1. By Lemma 2.3, it suffices to prove that  $X$  has a  $\sigma$ -hereditarily closure-preserving  $cs^*$ -network. Since  $f$  is an  $\sigma$ -map, there exists a base  $\mathcal{B}$  of  $Z$  such that  $f(\mathcal{B})$  is a  $\sigma$ -locally-finite collection in  $X$ , so we only need to prove that  $f(\mathcal{B})$  is a  $cs^*$ -network of  $X$ . Let  $S$  be a sequence converging to  $x \in X$  and  $U$  an open neighborhood of  $x$ .  $f$  is sequentially-quotient, so there exists a sequence  $L$  converging to  $z \in Z$  such that  $f(L)$  is a subsequence of  $S$ . Since  $z \in f^{-1}(x) \subset f^{-1}(U)$  and  $\mathcal{B}$  is a base of  $Z$ , there exists  $B \in \mathcal{B}$  such that  $z \in B \subset f^{-1}(U)$ . So  $L$  is eventually in  $B$ , hence  $f(L)$  is eventually in  $f(B) \subset ff^{-1}(U) = U$ . Thus  $S$  is frequently in  $f(B) \in f(\mathcal{B})$ . So  $f(\mathcal{B})$  is a  $cs^*$ -network of  $X$ .  $\square$

**COROLLARY 3.5.** *A space is  $g$ -metrizable if and only if it is a quotient,  $\pi$  (compact),  $\sigma$ -image of a metric space.*

**COROLLARY 3.6.** *Pseudo-open,  $\pi$ ,  $\sigma$ -maps preserve metric spaces.*

Recall that a map  $f : X \rightarrow Y$  is a  $\sigma$ -locally finite [15] if for every  $\sigma$ -locally finite cover  $\mathcal{P}$  of  $X$ , there exists a refinement  $\mathcal{B}$  of  $\mathcal{P}$  such that  $f(\mathcal{B})$  is a  $\sigma$ -locally finite collection. It is known that every  $\sigma$ -map is a  $\sigma$ -locally finite map, but the converse is not true [13]. We point out that “ $\sigma$ -map” in Theorem 3.4 can not be replaced

by “ $\sigma$ -locally-finite map”. In fact, the authors of [13] gave a compact-covering, open, compact,  $\sigma$ -locally finite map  $f$  from a metric space  $M$  onto a first countable non- $g$ -metrizable space  $X$  [13]. Then  $X$  is not  $sn$ -metrizable by Remark 1.5.

The author would like to thank the referee for his valuable amendments.

### References

- [1] P. S. Alexandroff and P. Urysohn, *Sur les espaces topologiques compacts*, Bull. Inter. Acad. Pol. Sci. A(1923), 5–8.
- [2] P. S. Alexandroff, *On some results concerning topological spaces and their continuous mappings*, Proc. 1st. Prague Top. Symp., 1961, 41–54.
- [3] A. Arhangel’skii, *Mappings and spaces*, Russian Math. Surveys 21(1966), 115–162.
- [4] J. R. Boone and F. Siwiec, *Sequentially quotient mappings*, Czech. Math. J. 26(1976) 174–182.
- [5] S. P. Franklin, *Spaces in which sequences suffice*, Fund. Math. 57(1965), 107–115.
- [6] J. A. Guthrie, *A characterization of  $\aleph_0$ -spaces*, Gen. Top. Appl. 1(1971), 105–110.
- [7] Y. Ge, *On spaces with a  $\sigma$ -locally-finite universal  $cs$ -network*, Questions Answers in General Topology, 18(2000), 93–96.
- [8] Y. Ge, *On  $sn$ -metrizable spaces*, Acta Math. Sinica 45(2002), 355–360 (in Chinese).
- [9] S. Lin, *Generalized metric spaces and mappings*, Chinese Science Press, Beijing, 1995 (in Chinese).
- [10] S. Lin, *A note on the Arens’ spaces and sequential fan*, Topology Appl. 81(1997), 185–196
- [11] S. Lin and P. Yan, *Sequence-covering maps of metric spaces*, Topology Appl. 109(2001), 301–314.
- [12] S. Lin and P. Yan, *On sequence-covering compact mappings*, Acta Math. Sinica 44(2001), 175–182 (in Chinese).
- [13] S. Lin and P. Yan, *Notes on  $cfp$ -covers*, Comment. Math. Univ. Carolinae 44(2003), 295–306.
- [14] E. A. Michael, *A quintuple quotient quest*, Gen. Top. Appl. 2(1972), 91–138.
- [15] E. A. Michael,  *$\sigma$ -locally finite maps*, Proc. Amer. Math. Soc. 65(1977), 159–164.
- [16] K. Morita, *A condition for the metrizability of topological spaces and for  $n$ -dimensionality*, Sci. Rep. Tokyo Kyoiku Daigaku Sec. A5(1955), 33–36.
- [17] V. I. Ponomarev, *Axioms of countability and continuous mappings*, Bull. Pol. Acad. Math. 8(1960), 127–133.
- [18] Y. Tanaka, *Symmetric spaces,  $g$ -developable spaces and  $g$ -metrizable spaces*, Math. Japonica 36(1991), 71–84.

Department of Mathematics  
Suzhou University, Suzhou 215006  
P. R. China  
geying@pub.sz.jsinfo.net

(Received 25 09 2003)